

Rabbit anti-Human, Mouse RELB Polyclonal Antibody | anti-RELB antibody

RELB Antibody

CATALOG # AAA233880	HOST Rabbit	REACTIVITY Human, Mouse	APPLICATIONS Immunofluorescence, Western Blot, ELISA
-------------------------------	-----------------------	-----------------------------------	--


PRODUCT SYNONYMS

RELB; Polyclonal Antibody; RELB Antibody; Transcription factor RelB; I-Rel; anti-RELB antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	579
Applicable Applications for anti-RELB antibody	IF (Immunofluorescence), WB (Western Blot), ELISA
Immunogen	Recombinant Human Transcription factor RelB protein (400-579AA)
Conjugation	Non-conjugated
Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IF (Immunofluorescence)

Immunofluorescent analysis of HeLa cells using AAA233880 at a dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)



WB (Western Blot)

Western BlotPositive WB detected in: NIH/3T3 whole cell lysateAll lanes: RELB antibody at 3ug/mlSecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 63, 62, 15 KDaObserved band size: 63 KDa

ADDITIONAL INFORMATION

Related Product Information for anti-RELB antibody	NF-kappa-B is a pleiotropic transcription factor which is present in almost all cell types and is involved in many biological processes such as inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NF-kappa-B is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NFKB1/p105, NFKB1/p50, REL and NFKB2/p52. The dimers bind at kappa-B sites in the DNA of their target genes and the individual dimers have distinct preferences for different kappa-B sites that they can bind with distinguishable affinity and specificity. Different dimer combinations act as transcriptional activators or repressors, respectively. NF-kappa-B is controlled by various mechanisms of post-translational modification and subcellular compartmentalization as well as by interactions with other cofactors or corepressors. NF-kappa-B complexes are held in the cytoplasm in an inactive state complexed with members of the NF-kappa-B inhibitor (I-kappa-B) family. In a conventional activation pathway, I-kappa-B is phosphorylated by I-kappa-B kinases (IKKs) in response to different activators, subsequently degraded thus liberating the active NF-kappa-B complex which translocates to the nucleus. NF-kappa-B heterodimeric RelB-p50 and RelB-p52 complexes are transcriptional activators. RELB neither associates with DNA nor with RELA/p65 or REL. Stimulates promoter activity in the presence of NFKB2/p49. As a member of the NUPR1/RELB/IER3 survival pathway, may provide pancreatic ductal adenocarcinoma with remarkable resistance to cell stress, such as starvation or gemcitabine treatment. Regulates the circadian clock by repressing the transcriptional activator activity of the CLOCK-ARNTL/BMAL1 heterodimer in a CRY1/CRY2 independent manner. Increased repression of the heterodimer is seen in the presence of NFKB2/p52.
Product Categories/Family for anti-RELB antibody	Epigenetics and Nuclear Signaling; Cancer; Cell biology; Signal transduction

NCBI AND UNIPROT INFORMATION

NCBI GI #	35493878
NCBI GeneID	5971
NCBI Accession #	NP_006500.2
NCBI GenBank Nucleotide #	NM_006509.3
UniProt Accession #	Q01201; Q6GTX7; Q9UEI7
Molecular Weight	62,134 Da
NCBI Official Full Name	transcription factor RelB
NCBI Official Synonym Full Names	RELB proto-oncogene, NF-kB subunit
NCBI Official Symbol	RELB
NCBI Official Synonym Symbols	IREL; I-REL; IMD53; REL-B
NCBI Protein Information	transcription factor RelB
UniProt Protein Name	Transcription factor RelB
UniProt Gene Name	RELB

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/reb-antibody-immunofluorescence-western-blot-elisa/233880>