

Rabbit anti-Human MLST8 Polyclonal Antibody | anti-MLST8 antibody

MLST8 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA233910	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

MLST8; Polyclonal Antibody; MLST8 Antibody; Target of rapamycin complex subunit LST8; TORC subunit LST8; G protein beta subunit-like; Gable; Protein GbetaL; Mammalian lethal with SEC13 protein 8; mLST8; GBL; LST8; anti-MLST8 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	326
Applicable Applications for anti-MLST8 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant Human Target of rapamycin complex subunit LST8 protein (159-220AA)
Conjugation	Non-conjugated
Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IF (Immunofluorescence)

Immunofluorescent analysis of MCF-7 cells using AAA233910 at a dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human kidney tissue using AAA233910 at dilution of 1:100

ADDITIONAL INFORMATION

Related Product Information for anti-MLST8 antibody	Subunit of both mTORC1 and mTORC2, which regulates cell growth and survival in response to nutrient and hormonal signals. mTORC1 is activated in response to growth factors or amino acids. Growth factor-stimulated mTORC1 activation involves a AKT1-mediated phosphorylation of TSC1-TSC2, which leads to the activation of the RHEB GTPase that potently activates the protein kinase activity of mTORC1. Amino acid-signaling to mTORC1 requires its relocalization to the lysosomes mediated by the Ragulator complex and the Rag GTPases. Activated mTORC1 up-regulates protein synthesis by phosphorylating key regulators of mRNA translation and ribosome synthesis. mTORC1 phosphorylates EIF4EBP1 and releases it from inhibiting the elongation initiation factor 4E (eiF4E). mTORC1 phosphorylates and activates S6K1 at 'Thr-389', which then promotes protein synthesis by phosphorylating PDCD4 and targeting it for degradation. Within mTORC1, LST8 interacts directly with MTOR and enhances its kinase activity. In nutrient-poor conditions, stabilizes the MTOR-RPTOR interaction and favors RPTOR-mediated inhibition of MTOR activity. mTORC2 is also activated by growth factors, but seems to be nutrient-insensitive. mTORC2 seems to function upstream of Rho GTPases to regulate the actin cytoskeleton, probably by activating one or more Rho-type guanine nucleotide exchange factors. mTORC2 promotes the serum-induced formation of stress-fibers or F-actin. mTORC2 plays a critical role in AKT1 'Ser-473' phosphorylation, which may facilitate the phosphorylation of the activation loop of AKT1 on 'Thr-308' by PDK1 which is a prerequisite for full activation. mTORC2 regulates the phosphorylation of SGK1 at 'Ser-422'. mTORC2 also modulates the phosphorylation of PRKCA on 'Ser-657'.
Product Categories/Family for anti-MLST8 antibody	Epigenetics and Nuclear Signaling; Signal Transduction

NCBI AND UNIPROT INFORMATION

NCBI GI #	312596910
NCBI GeneID	64223
NCBI Accession #	NP_001186102.1
NCBI GenBank Nucleotide #	NM_001199173.2
UniProt Accession #	Q9BVC4; Q5M800; Q86Y18; Q8WUI5; Q9HA66; Q9UJV6; B3KMM4; B4DY00; D3DU88
Molecular Weight	35,748 Da
NCBI Official Full Name	target of rapamycin complex subunit LST8 isoform a
NCBI Official Synonym Full Names	MTOR associated protein, LST8 homolog
NCBI Official Symbol	MLST8
NCBI Official Synonym Symbols	GBL; LST8; POP3; WAT1; GbetaL
NCBI Protein Information	target of rapamycin complex subunit LST8
UniProt Protein Name	Target of rapamycin complex subunit LST8
UniProt Gene Name	MLST8
UniProt Synonym Gene Names	GBL; LST8; TORC subunit LST8; Gable; Protein GbetaL; mLST8

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/mlst8-antibody-immunofluorescence-immunohistochemistry-elisa/233910>