

Rabbit anti-Human GDI1 Polyclonal Antibody | anti-GDI1 antibody

GDI1 Antibody

CATALOG # AAA234076	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

GDI1; Polyclonal Antibody; GDI1 Antibody; Rab GDP dissociation inhibitor alpha; Rab GDI alpha; Guanosine diphosphate dissociation inhibitor 1; GDI-1; Oligophrenin-2; Protein XAP-4; GDIL; OPHN2; RABGDIA; XAP4; anti-GDI1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	447
Applicable Applications for anti-GDI1 antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Recombinant Human Rab GDP dissociation inhibitor alpha protein (35-170AA)
Conjugation	Non-conjugated
Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IHC (Immunohistochemistry)

IHC image of AAA234076 diluted at 1:200 and staining in paraffin-embedded human prostate cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)

IHC image of AAA234076 diluted at 1:200 and staining in paraffin-embedded human brain tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



WB (Western Blot)

Western Blot Positive WB detected in: A549 whole cell lysate
 All lanes: GDI1 antibody at 6 µg/ml
 Secondary: Goat polyclonal to rabbit IgG at 1/50000 dilution
 Predicted band size: 51 KDa
 Observed band size: 51 KDa

ADDITIONAL INFORMATION

Related Product Information for anti-GDI1 antibody	Regulates the GDP/GTP exchange reaction of most Rab proteins by inhibiting the dissociation of GDP from them, and the subsequent binding of GTP to them. Promotes the dissociation of GDP-bound Rab proteins from the membrane and inhibits their activation. Promotes the dissociation of RAB1A, RAB3A, RAB5A and RAB10 from membranes.
Product Categories/Family for anti-GDI1 antibody	Signal Transduction

NCBI AND UNIPROT INFORMATION

NCBI GI #	4503971
NCBI GeneID	2664
NCBI Accession #	NP_001484.1
NCBI GenBank Nucleotide #	NM_001493.2
UniProt Accession #	P31150; P50394; Q6FG50; Q7Z2G6; Q7Z2G9; Q7Z2H5; Q7Z2I6
Molecular Weight	50,583 Da
NCBI Official Full Name	rab GDP dissociation inhibitor alpha
NCBI Official Synonym Full Names	GDP dissociation inhibitor 1
NCBI Official Symbol	GDI1
NCBI Official Synonym Symbols	1A; GDIL; MRX41; MRX48; OPHN2; XAP-4; RABGD1A; RABGDIA
NCBI Protein Information	rab GDP dissociation inhibitor alpha
UniProt Protein Name	Rab GDP dissociation inhibitor alpha
UniProt Gene Name	GDI1
UniProt Synonym Gene Names	GDIL; OPHN2; RABGDIA; XAP4; Rab GDI alpha; GDI-1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/gdi1-antibody-immunohistochemistry-western-blot-elisa/234076>