

Rabbit anti-Human, Rat MCM4 Polyclonal Antibody | anti-MCM4 antibody

MCM4 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA234084	Rabbit	Human, Rat	Immunofluorescence, Immunohistochemistry, Western Blot, ELISA


PRODUCT SYNONYMS

MCM4; Polyclonal Antibody; MCM4 Antibody; DNA replication licensing factor MCM4; CDC21 homolog; P1-CDC21; CDC21; anti-MCM4 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	863
Applicable Applications for anti-MCM4 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Recombinant Human DNA replication licensing factor MCM4 protein (2-141AA)
Conjugation	Non-conjugated
Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IHC (Immunohistochemistry)

Immunofluorescence staining of A549 cells with AAA234084 at 1:133, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L).



IHC (Immunohistochemistry)

IHC image of AAA234084 diluted at 1:100 and staining in paraffin-embedded human testis tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



WB (Western Blot)

Western Blot Positive WB detected in: HeLa whole cell lysate, 293 whole cell lysate, Rat heart tissue
 All lanes: MCM4 antibody at 2.7ug/ml Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution
 Predicted band size: 97 KDa Observed band size: 97 KDa

ADDITIONAL INFORMATION

Related Product Information for anti-MCM4 antibody	Acts as component of the MCM2-7 complex (MCM complex) which is the putative replicative helicase essential for 'once per cell cycle' DNA replication initiation and elongation in eukaryotic cells. The active ATPase sites in the MCM2-7 ring are formed through the interaction surfaces of two neighboring subunits such that a critical structure of a conserved arginine finger motif is provided in trans relative to the ATP-binding site of the Walker A box of the adjacent subunit. The six ATPase active sites, however, are likely to contribute differentially to the complex helicase activity.
Product Categories/Family for anti-MCM4 antibody	Epigenetics and Nuclear Signaling

NCBI AND UNIPROT INFORMATION

NCBI GI #	33469919
NCBI GeneID	4173
NCBI Accession #	NP_005905.2
NCBI GenBank Nucleotide #	NM_005914.3
UniProt Accession #	P33991 ; Q8NEH1 ; Q99658
Molecular Weight	96,558 Da
NCBI Official Full Name	DNA replication licensing factor MCM4
NCBI Official Synonym Full Names	minichromosome maintenance complex component 4
NCBI Official Symbol	MCM4
NCBI Official Synonym Symbols	NKCD; CDC21; CDC54; IMD54; NKGCD; hCdc21; P1-CDC21
NCBI Protein Information	DNA replication licensing factor MCM4
UniProt Protein Name	DNA replication licensing factor MCM4
UniProt Gene Name	MCM4
UniProt Synonym Gene Names	CDC21

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.