

Rabbit anti-Human EPHA2 Polyclonal Antibody | anti-EPHA2 antibody

EPHA2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA234203	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

EPHA2; Polyclonal Antibody; EPHA2 Antibody; Ephrin type-A receptor 2; Epithelial cell kinase; Tyrosine-protein kinase receptor ECK; ECK; anti-EPHA2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	976
Applicable Applications for anti-EPHA2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant Human Ephrin type-A receptor 2 protein (627-976AA)
Conjugation	Non-conjugated
Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IF (Immunofluorescence)

Immunofluorescence staining of A549 cells with AAA234203 at 1:133, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L).



IHC (Immunohistochemistry)

IHC image of AAA234203 diluted at 1:400 and staining in paraffin-embedded human brain tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

ADDITIONAL INFORMATION

Related Product Information for anti-EPHA2 antibody	Receptor tyrosine kinase which binds promiscuously membrane-bound ephrin-A family ligands residing on adjacent cells, leading to contact-dependent bidirectional signaling into neighboring cells. The signaling pathway downstream of the receptor is referred to as forward signaling while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling. Activated by the ligand ephrin-A1/EFNA1 regulates migration, integrin-mediated adhesion, proliferation and differentiation of cells. Regulates cell adhesion and differentiation through DSG1/desmoglein-1 and inhibition of the ERK1/ERK2 (MAPK3/MAPK1, respectively) signaling pathway. May also participate in UV radiation-induced apoptosis and have a ligand-independent stimulatory effect on chemotactic cell migration. During development, may function in distinctive aspects of pattern formation and subsequently in development of several fetal tissues. Involved for instance in angiogenesis, in early hindbrain development and epithelial proliferation and branching morphogenesis during mammary gland development. Engaged by the ligand ephrin-A5/EFNA5 may regulate lens fiber cells shape and interactions and be important for lens transparency development and maintenance. With ephrin-A2/EFNA2 may play a role in bone remodeling through regulation of osteoclastogenesis and osteoblastogenesis.
Product Categories/Family for anti-EPHA2 antibody	Neuroscience; Cardiovascular

NCBI AND UNIPROT INFORMATION

NCBI GI #	32967311
NCBI GeneID	1969
NCBI Accession #	NP_004422.2
NCBI GenBank Nucleotide #	NM_004431.4
UniProt Accession #	P29317; Q8N3Z2; B5A968
Molecular Weight	54,305 Da
NCBI Official Full Name	ephrin type-A receptor 2 isoform 1
NCBI Official Synonym Full Names	EPH receptor A2
NCBI Official Symbol	EPHA2
NCBI Official Synonym Symbols	ECK; CTPA; ARCC2; CTPP1; CTRCT6
NCBI Protein Information	ephrin type-A receptor 2
UniProt Protein Name	Ephrin type-A receptor 2
UniProt Gene Name	EPHA2
UniProt Synonym Gene Names	ECK

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/epha2-antibody-immunofluorescence-immunohistochemistry-elisa/234203>