

Rabbit anti-Human, Mouse HIST1H2AG Polyclonal Antibody | anti-HIST1H2AG antibody

HIST1H2AG (Ab-36) Antibody

CATALOG # AAA234372	HOST Rabbit	REACTIVITY Human, Mouse	APPLICATIONS Immunofluorescence, Western Blot, ELISA
-------------------------------	-----------------------	-----------------------------------	--


PRODUCT SYNONYMS

HIST1H2AG; Polyclonal Antibody; HIST1H2AG (Ab-36) Antibody; Histone H2A type 1; H2A.1; Histone H2A/p; H2AFP; AND; HIST1H2AI; H2AFC; HIST1H2AK; H2AFD; HIST1H2AL; H2AFI; HIST1H2AM; H2AFN; anti-HIST1H2AG antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid
Sequence Length	130
Applicable Applications for anti-HIST1H2AG antibody	IF (Immunofluorescence), WB (Western Blot), ELISA
Immunogen	Peptide sequence around site of Lys (36) derived from human Histone H2A type 1.
Conjugation	Non-conjugated
Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IF (Immunofluorescence)

Immunofluorescence staining of hela with AAA234372, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L).



WB (Western Blot)

Western BlotPositive WB detected in: Mouse heart tissueAll lanes: HIST1H2AG antibody at 1.04ug/mlSecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 15 KDaObserved band size: 15 KDa

ADDITIONAL INFORMATION

Related Product Information for anti-HIST1H2AG antibody	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
Product Categories/Family for anti-HIST1H2AG antibody	Epigenetics and Nuclear Signaling

NCBI AND UNIPROT INFORMATION

NCBI GI #	4504239
NCBI GeneID	8329
NCBI Accession #	NP_003500.1
NCBI GenBank Nucleotide #	NM_003509.2
UniProt Accession #	P0C0S8; P02261; Q2M1R2; Q76PA6
Molecular Weight	14,091 Da
NCBI Official Full Name	histone H2A type 1
NCBI Official Synonym Full Names	histone cluster 1 H2A family member i
NCBI Official Symbol	HIST1H2AI
NCBI Official Synonym Symbols	H2A/c; H2AFC
NCBI Protein Information	histone H2A type 1
UniProt Protein Name	Histone H2A type 1
UniProt Gene Name	HIST1H2AG
UniProt Synonym Gene Names	H2AFP; H2A.1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/hist1h2ag-antibody-immunofluorescence-western-blot-elisa/234372>