

Rabbit anti-Human SKA3 Polyclonal Antibody | anti-SKA3 antibody

SKA3 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA234481	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

SKA3; Polyclonal Antibody; SKA3 Antibody; Spindle and kinetochore-associated protein 3; C13orf3; RAMA1; anti-SKA3 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	388
Applicable Applications for anti-SKA3 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant Human Spindle and kinetochore-associated protein 3 protein (101-195AA)
Conjugation	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Immunogen Species	Homo sapiens (Human)
Research Area	Cell Biology
Target Names	SKA3
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.


IMAGES



IHC (Immunohistochemisry)
Immunofluorescence staining of Hela cells with AAA234481 at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L).



IHC (Immunohiostchemistry)
IHC image of AAA234481 diluted at 1:300 and staining in paraffin-embedded human melanoma cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)
IHC image of AAA234481 diluted at 1:300 and staining in paraffin-embedded human testis tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

ADDITIONAL INFORMATION

Product Categories/Family for anti-SKA3 antibody	Cell Biology
--	--------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	260763912
NCBI GeneID	221150
NCBI Accession #	NP_001159489.1
NCBI GenBank Nucleotide #	NM_001166017.1
UniProt Accession #	Q8IX90; Q5VZV5; Q86WR2; Q8NBG1; Q96D22; A2A330; A2A331; B2RBY2
NCBI Official Full Name	spindle and kinetochore-associated protein 3 isoform 2
NCBI Official Synonym Full Names	spindle and kinetochore associated complex subunit 3
NCBI Official Symbol	SKA3
NCBI Official Synonym Symbols	RAMA1; C13orf3

NCBI Protein Information	spindle and kinetochore-associated protein 3
UniProt Protein Name	Spindle and kinetochore-associated protein 3
UniProt Gene Name	SKA3
UniProt Synonym Gene Names	C13orf3; RAMA1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/ska3-antibody-immunofluorescence-immunohistochemistry-elisa/234481>