

Rabbit anti-Human HIST1H3A Polyclonal Antibody | anti-HIST1H3A antibody

HIST1H3A(Ab-64) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA234539	Rabbit	Human	Immunohistochemistry, ELISA


PRODUCT SYNONYMS

HIST1H3A; Polyclonal Antibody; HIST1H3A(Ab-64) Antibody; Histone H3.1; Histone H3/a; Histone H3/b; Histone H3/c; Histone H3/d; Histone H3/f; Histone H3/h; Histone H3/i; Histone H3/j; Histone H3/k; Histone H3/l; H3FA; AND; HIST1H3B; H3FL; HIST1H3C; H3FC; HIST1H3D; H3FB; HIST1H3E; H3FD; HIST1H3F; H3FI; HIST1H3G; H3FH; HIST1H3H; H3FK; HIST1H3I; H3FF; HIST1H3J; H3FJ; anti-HIST1H3A antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid
Sequence Length	136
Applicable Applications for anti-HIST1H3A antibody	IHC (Immunohistochemistry), ELISA
Immunogen	Peptide sequence around site of acetyl-Lys(64) derived from human Histone H3
Conjugation	Non-conjugated
Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)
IHC image of AAA234539 diluted at 1:100 and staining in paraffin-embedded human bladder cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)
IHC image of AAA234539 diluted at 1:100 and staining in paraffin-embedded human prostate cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

ADDITIONAL INFORMATION

Product Categories/Family for anti-HIST1H3A antibody	Epigenetics and Nuclear Signaling
--	-----------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	4504281
NCBI GeneID	8350
NCBI Accession #	NP_003520.1
NCBI GenBank Nucleotide #	NM_003529.2
UniProt Accession #	P68431; P02295; P02296; P16106; Q6ISV8; Q6NWP8; Q6NWP9; Q6NXU4; Q71DJ3; Q93081; A0PJT7; A5PLR1
Molecular Weight	15,404 Da
NCBI Official Full Name	histone H3.1
NCBI Official Synonym Full Names	histone cluster 1 H3 family member a
NCBI Official Symbol	HIST1H3A
NCBI Official Synonym Symbols	H3/A; H3FA
NCBI Protein Information	histone H3.1
UniProt Protein Name	Histone H3.1
UniProt Gene Name	HIST1H3A
UniProt Synonym Gene Names	H3FA

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.