

Rabbit HIST1H4A Polyclonal Antibody | anti-HIST1H4A antibody

Butyryl-HIST1H4A (K5) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA234739	Rabbit	Human, Mouse, Rat	Immunofluorescence, Western Blot, ELISA


PRODUCT SYNONYMS

HIST1H4A; Polyclonal Antibody; Butyryl-HIST1H4A (K5) Antibody; Histone H4; H4/A; H4FA; AND; HIST1H4B; H4/I; H4FI; HIST1H4C; H4/G; H4FG; HIST1H4D; H4/B; H4FB; HIST1H4E; H4/J; H4FJ; HIST1H4F; H4/C; H4FC; HIST1H4H; H4/H; H4FH; HIST1H4I; H4/M; H4FM; HIST1H4J; H4/E; H4FE; HIST1H4K; H4/D; H4FD; HIST1H4L; H4/K; H4FK; HIST2H4A; H4/N; H4F2; H4FN; HIST2H4; HIST2H4B; H4/O; H4FO; HIST4H4; anti-HIST1H4A antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid
Sequence Length	103
Applicable Applications for anti-HIST1H4A antibody	IF (Immunofluorescence), WB (Western Blot), ELISA
Immunogen	Peptide sequence around site of butyryl-Lys(5) derived from human Histone H4
Conjugation	Non-conjugated
Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IF (Immunofluorescence)
Immunofluorescence staining of HepG2 cells(treated with 30mM sodium butyrate for 4h) with AAA234739, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L).



WB (Western Blot)
Western BlotPositive WB detected in: HeLa whole cell lysate, Jurkat whole cell lysate, 293 whole cell lysate, HepG2 whole cell lysate (treated with 30mM sodium butyrate for 4h)All lanes: HIST1H4A antibody at 0.14ug/mlSecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 12 KDaObserved band size: 12 KDa

ADDITIONAL INFORMATION

Related Product Information for anti-HIST1H4A antibody	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
Product Categories/Family for anti-HIST1H4A antibody	Epigenetics and Nuclear Signaling

NCBI AND UNIPROT INFORMATION

NCBI GI #	77539758
NCBI GeneID	554313
NCBI Accession #	NP_001029249.1
NCBI GenBank Nucleotide #	NM_001034077.4
UniProt Accession #	P62805; P02304; P02305; Q6DRA9; Q6FGB8; Q6NWP7; A2VCL0
Molecular Weight	11,367 Da
NCBI Official Full Name	histone H4
NCBI Official Synonym Full Names	histone cluster 2 H4 family member b
NCBI Official Symbol	HIST2H4B
NCBI Official Synonym Symbols	H4/o
NCBI Protein Information	histone H4
UniProt Protein Name	Histone H4
UniProt Gene Name	HIST1H4A
UniProt Synonym Gene Names	H4/A; H4FA

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

