

Rabbit anti-Human HIST1H4A Polyclonal Antibody | anti-HIST1H4A antibody

HIST1H4A(Ab-44) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA234924	Rabbit	Human	Immunohistochemistry, ELISA


PRODUCT SYNONYMS

HIST1H4A; Polyclonal Antibody; HIST1H4A(Ab-44) Antibody; Histone H4; H4/A; H4FA; AND; HIST1H4B; H4/I; H4FI; HIST1H4C; H4/G; H4FG; HIST1H4D; H4/B; H4FB; HIST1H4E; H4/J; H4FJ; HIST1H4F; H4/C; H4FC; HIST1H4H; H4/H; H4FH; HIST1H4I; H4/M; H4FM; HIST1H4J; H4/E; H4FE; HIST1H4K; H4/D; H4FD; HIST1H4L; H4/K; H4FK; HIST2H4A; H4/N; H4F2; H4FN; HIST2H4; HIST2H4B; H4/O; H4FO; HIST4H4; anti-HIST1H4A antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid
Sequence Length	103
Applicable Applications for anti-HIST1H4A antibody	IHC (Immunohistochemistry), ELISA
Immunogen	Peptide sequence around site of butyryl-Lys(44) derived from human Histone H4
Conjugation	Non-conjugated
Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)
IHC image of AAA234924 diluted at 1:15 and staining in paraffin-embedded human lung cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)
IHC image of AAA234924 diluted at 1:15 and staining in paraffin-embedded human liver cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

ADDITIONAL INFORMATION

Product Categories/Family for anti-HIST1H4A antibody	Epigenetics and Nuclear Signaling
--	-----------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	77539758
NCBI GeneID	554313
NCBI Accession #	NP_001029249.1
NCBI GenBank Nucleotide #	NM_001034077.4
UniProt Accession #	P62805; P02304; P02305; Q6DRA9; Q6FGB8; Q6NWP7; A2VCL0
Molecular Weight	11,367 Da
NCBI Official Full Name	histone H4
NCBI Official Synonym Full Names	histone cluster 2 H4 family member b
NCBI Official Symbol	HIST2H4B
NCBI Official Synonym Symbols	H4/o
NCBI Protein Information	histone H4
UniProt Protein Name	Histone H4
UniProt Gene Name	HIST1H4A
UniProt Synonym Gene Names	H4/A; H4FA

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.