

Rabbit anti-Human TCF20 Polyclonal Antibody | anti-TCF20 antibody

TCF20 Antibody

CATALOG # AAA235052	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

TCF20; Polyclonal Antibody; TCF20 Antibody; Transcription factor 20; TCF-20; Nuclear factor SPBP; Protein AR1; Stromelysin-1 PDGF-responsive element-binding protein; SPRE-binding protein; KIAA0292; SPBP; anti-TCF20 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	1960
Applicable Applications for anti-TCF20 antibody	IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant Human Transcription factor 20 protein (539-704AA)
Conjugation	Non-conjugated
Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)

IHC image of AAA235052 diluted at 1:300 and staining in paraffin-embedded human glioma cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)

IHC image of AAA235052 diluted at 1:300 and staining in paraffin-embedded human brain tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

ADDITIONAL INFORMATION

Related Product Information for anti-TCF20 antibody	Transcriptional activator that binds to the regulatory region of MMP3 and thereby controls stromelysin expression. It stimulates the activity of various transcriptional activators such as JUN, SP1, PAX6 and ETS1, suggesting a function as a coactivator.
Product Categories/Family for anti-TCF20 antibody	Epigenetics and Nuclear Signaling

NCBI AND UNIPROT INFORMATION

NCBI GI #	31652244
NCBI GeneID	6942
NCBI Accession #	NP_005641.1
NCBI GenBank Nucleotide #	NM_005650.3
UniProt Accession #	Q9UGU0; O14528; Q13078; Q4V353; Q9H4M0; A9JX12
Molecular Weight	209,654 Da
NCBI Official Full Name	transcription factor 20 isoform 1
NCBI Official Synonym Full Names	transcription factor 20
NCBI Official Symbol	TCF20
NCBI Official Synonym Symbols	AR1; SPBP; TCF-20
NCBI Protein Information	transcription factor 20
UniProt Protein Name	Transcription factor 20
UniProt Gene Name	TCF20
UniProt Synonym Gene Names	KIAA0292; SPBP; TCF-20; SPRE-binding protein

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/tcf20-antibody-immunohistochemistry-elisa/235052>