

Rabbit anti-Human TCAF2 Polyclonal Antibody | anti-TCAF2 antibody

TCAF2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA235297	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

TCAF2; Polyclonal Antibody; TCAF2 Antibody; TRPM8 channel-associated factor 2; TRP channel-associated factor 2; FAM115C FAM139A; anti-TCAF2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95% Protein G Purified
Form/Format	Liquid
Sequence Length	845
Applicable Applications for anti-TCAF2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen Species	Human
Immunogen	Recombinant Human TRPM8 channel-associated factor 2 protein (472-590AA)
Conjugate	Non-Conjugated
Preservative	0.03% Proclin 300
Constituents	50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.


IMAGES



IF (Immunofluorescence)
Immunofluorescence staining of HeLa cells with CSB-PA008015LA01HU at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IHC (Immunohistochemistry)
IHC image of CSB-PA008015LA01HU diluted at 1:300 and staining in paraffin-embedded human liver cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)
IHC image of CSB-PA008015LA01HU diluted at 1:300 and staining in paraffin-embedded human pancreatic cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

ADDITIONAL INFORMATION

Product Categories/Family for anti-TCAF2 antibody	Cell Biology
---	--------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	194097428
NCBI GeneID	285966
NCBI Accession #	NP_001123497.1
NCBI GenBank Nucleotide #	NM_001130025.1
UniProt Accession #	A6NFQ2; Q14D25; Q17RQ4; Q8IWQ0; Q8NF84; B4DK02
Molecular Weight	92,008 Da
NCBI Official Synonym Full Names	TRPM8 channel associated factor 2
NCBI Official Symbol	TCAF2
NCBI Official Synonym Symbols	GATD9; FAM115C; FAM139A

NCBI Protein Information	TRPM8 channel-associated factor 2
UniProt Protein Name	TRPM8 channel-associated factor 2
UniProt Gene Name	TCAF2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/tcaf2-antibody-immunofluorescence-immunohistochemistry-elisa/235297>