

Rabbit anti-Human SKOR2 Polyclonal Antibody | anti-SKOR2 antibody

SKOR2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA235305	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

SKOR2; Polyclonal Antibody; SKOR2 Antibody; SKI family transcriptional corepressor 2; Functional Smad-suppressing element on chromosome 18; Fussel-18; LBX1 corepressor 1-like protein; Ladybird homeobox corepressor 1-like protein; CORL2 FUSSEL18; anti-SKOR2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Sequence Length	296
Applicable Applications for anti-SKOR2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen Species	Human
Immunogen	Recombinant Human SKI family transcriptional corepressor 2 protein (19-36AA)
Conjugate	Non-Conjugated
Research Area	Signal Transduction
Target Names	SKOR2
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IF (Immunofluorescence)

Immunofluorescence staining of K562 cells with AAA235305 at 1:50, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IHC (Immunohistochemistry)

IHC image of AAA235305 diluted at 1:100 and staining in paraffin-embedded human colon cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)

IHC image of AAA235305 diluted at 1:100 and staining in paraffin-embedded human prostate cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

ADDITIONAL INFORMATION

Product Categories/Family for anti-SKOR2 antibody	Signal Transduction
---	---------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	83699422
NCBI GeneID	652991
NCBI Accession #	NP_001032891.1
NCBI GenBank Nucleotide #	NM_001037802.1
UniProt Accession #	Q2VWA4
NCBI Official Full Name	SKI family transcriptional corepressor 2 isoform 2
NCBI Official Synonym Full Names	SKI family transcriptional corepressor 2
NCBI Official Symbol	SKOR2
NCBI Official Synonym Symbols	CORL2; CH18515; FUSSEL18

NCBI Protein Information	SKI family transcriptional corepressor 2
UniProt Protein Name	SKI family transcriptional corepressor 2
UniProt Gene Name	SKOR2
UniProt Synonym Gene Names	CORL2; FUSSEL18; Fussel-18

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/skor2-antibody-immunofluorescence-immunohistochemistry-elisa/235305>