

Rabbit anti-Human KRT16 Polyclonal Antibody | anti-KRT16 antibody

KRT16 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA235325	Rabbit	Human	Immunofluorescence, Immunohistochemistry, Western Blot, ELISA


PRODUCT SYNONYMS

KRT16; Polyclonal Antibody; KRT16 Antibody; Keratin; type I cytoskeletal 16; Cytokeratin-16; CK-16; Keratin-16; K16; KRT16A; anti-KRT16 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95% Protein G Purified
Form/Format	Liquid
Sequence Length	473
Applicable Applications for anti-KRT16 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen Species	Human
Immunogen	Recombinant Human Keratin, type I cytoskeletal 16 protein (265-333AA)
Conjugate	Non-Conjugated
Preservative	0.03% Proclin 300
Constituents	50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.


IMAGES




IF (Immunofluorescence)
Immunofluorescence staining of MCF-7 cells with CSB-PA012515LA01HU at 1:133, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IHC (Immunohistochemistry)
IHC image of CSB-PA012515LA01HU diluted at 1:400 and staining in paraffin-embedded human tonsil tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)
IHC image of CSB-PA012515LA01HU diluted at 1:400 and staining in paraffin-embedded human cervical cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



WB (Western Blot)
Western BlotPositive WB detected in: HepG2 whole cell lysate, A549 whole cell lysateAll lanes: KRT16 antibody at 3.4ug/mlSecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 52 kDaObserved band size: 52 kDa

ADDITIONAL INFORMATION

Product Categories/Family for anti-KRT16 antibody	Signal Transduction
---	---------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	24430192
NCBI GeneID	3868
NCBI Accession #	NP_005548.2
NCBI GenBank Nucleotide #	NM_005557.4
UniProt Accession #	P08779; P30654; Q16402; Q9UBG8; A8K488
Molecular Weight	51,268 Da
NCBI Official Full Name	keratin, type I cytoskeletal 16
NCBI Official Synonym Full Names	keratin 16
NCBI Official Symbol	KRT16

NCBI Official Synonym Symbols	K16; PC1; CK16; K1CP; NEPPK; FNEPPK; KRT16A
NCBI Protein Information	keratin, type I cytoskeletal 16
UniProt Protein Name	Keratin, type I cytoskeletal 16
UniProt Gene Name	KRT16
UniProt Synonym Gene Names	KRT16A; CK-16; K16

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/krt16-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/235325>