

Rabbit anti-Human LIMD2 Polyclonal Antibody | anti-LIMD2 antibody

LIMD2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA235329	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

LIMD2; Polyclonal Antibody; LIMD2 Antibody; LIM domain-containing protein 2; anti-LIMD2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95% Protein G Purified
Form/Format	Liquid
Sequence Length	127
Applicable Applications for anti-LIMD2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen Species	Human
Immunogen	Recombinant Human LIM domain-containing protein 2 protein (1-127AA)
Conjugate	Non-Conjugated
Preservative	0.03% Proclin 300
Constituents	50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IF (Immunofluorescence)

Immunofluorescence staining of SH-SY5Y cells with CSB-PA012950LA01HU at 1:167, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IHC (Immunohistochemistry)

IHC image of CSB-PA012950LA01HU diluted at 1:500 and staining in paraffin-embedded human brain tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)

IHC image of CSB-PA012950LA01HU diluted at 1:500 and staining in paraffin-embedded human liver cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

ADDITIONAL INFORMATION

Product Categories/Family for anti-LIMD2 antibody	Cell Biology
---	--------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	13386490
NCBI GeneID	80774
NCBI Accession #	NP_085053.1
NCBI GenBank Nucleotide #	NM_030576.4
UniProt Accession #	Q9BT23; Q96S91; D3DU16
Molecular Weight	14,070 Da
NCBI Official Full Name	LIM domain-containing protein 2
NCBI Official Synonym Full Names	LIM domain containing 2
NCBI Official Symbol	LIMD2

NCBI Protein Information	LIM domain-containing protein 2
UniProt Protein Name	LIM domain-containing protein 2
UniProt Gene Name	LIMD2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/limd2-antibody-immunofluorescence-immunohistochemistry-elisa/235329>