

Rabbit anti-Human BCL2L10 Polyclonal Antibody | anti-BCL2L10 antibody

BCL2L10 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA235505	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

BCL2L10; Polyclonal Antibody; BCL2L10 Antibody; Bcl-2-like protein 10; Bcl2-L-10; Anti-apoptotic protein NrH; Apoptosis regulator Bcl-B; BCLB; anti-BCL2L10 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95% Protein G Purified
Form/Format	Liquid
Sequence Length	241
Applicable Applications for anti-BCL2L10 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen Species	Human
Immunogen	Recombinant Human Bcl-2-like protein 10 protein (19-127AA)
Conjugate	Non-Conjugated
Preservative	0.03% Proclin 300
Constituents	50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.


IMAGES



IF (Immunofluorescence)
Immunofluorescence staining of A549 cells with CSB-PA884627LA01HU at 1:50, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IHC (Immunohistochemistry)
IHC image of CSB-PA884627LA01HU diluted at 1:150 and staining in paraffin-embedded human liver cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)
IHC image of CSB-PA884627LA01HU diluted at 1:150 and staining in paraffin-embedded human kidney tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

ADDITIONAL INFORMATION

Product Categories/Family for anti-BCL2L10 antibody	Cell Biology
---	--------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	807201162
NCBI GeneID	10017
NCBI Accession #	NP_001293097.1
NCBI GenBank Nucleotide #	NM_001306168.1
UniProt Accession #	Q9HD36; Q3SX80; Q52LQ9; Q8TCS9
Molecular Weight	21,973 Da
NCBI Official Full Name	bcl-2-like protein 10 isoform 1
NCBI Official Synonym Full Names	BCL2 like 10
NCBI Official Symbol	BCL2L10

NCBI Official Synonym Symbols	Boo; Diva; BCL-B; bcl2-L-10
NCBI Protein Information	bcl-2-like protein 10
UniProt Protein Name	Bcl-2-like protein 10
UniProt Gene Name	BCL2L10
UniProt Synonym Gene Names	BCLB; Bcl2-L-10

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/bcl2l10-antibody-immunofluorescence-immunohistochemistry-elisa/235505>