

EIF4E recombinant antibody

Phospho-EIF4E (S209) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA235538	Rabbit	Human	Immunohistochemistry, Western Blot, ELISA


PRODUCT SYNONYMS

EIF4E; Monoclonal Recombinant Antibody; Phospho-EIF4E (S209) Antibody; Eukaryotic translation initiation factor 4E; eIF-4E; eIF4E; eIF-4F 25 kDa subunit; mRNA cap-binding protein; EIF4EL1; EIF4F; EIF4E recombinant antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG
Purity/Purification	Affinity-Chromatography
Form/Format	Liquid; In phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Sequence Length	237
Applicable Applications for EIF4E recombinant antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen Species	Human
Immunogen	A synthesized peptide derived from human Phospho-EIF4E (S209)
Conjugate	Non-Conjugated
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)

IHC image of CSB-RA007556A209pH1U diluted at 1:100 and staining in paraffin-embedded human breast cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



WB (Western Blot)

Western Blot Positive WB detected in 293 whole cell lysate, HepG2 whole cell lysate (treated with Calyculin A or not). All lanes? Phospho-EIF4E antibody at 1.2ug/ml. Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 25 kDa. Observed band size: 25 kDa.

ADDITIONAL INFORMATION

Related Product Information for EIF4E recombinant antibody	Recognizes and binds the 7-methylguanosine-containing mRNA cap during an early step in the initiation of protein synthesis and facilitates ribosome binding by inducing the unwinding of the mRNAs secondary structures. Component of the CYFIP1-EIF4E-FMR1 complex which binds to the mRNA cap and mediates translational repression. In the CYFIP1-EIF4E-FMR1 complex this subunit mediates the binding to the mRNA cap.
Product Categories/Family for EIF4E recombinant antibody	Recombinant Antibody; Epigenetics and Nuclear Signaling

NCBI AND UNIPROT INFORMATION

NCBI GI #	194578907
NCBI GeneID	1977
NCBI Accession #	NP_001124150.1
NCBI GenBank Nucleotide #	NM_001130678.2
UniProt Accession #	P06730; Q96E95; B7Z6V1; D6RCQ6
Molecular Weight	28,778 Da
NCBI Official Full Name	eukaryotic translation initiation factor 4E isoform 3
NCBI Official Synonym Full Names	eukaryotic translation initiation factor 4E
NCBI Official Symbol	EIF4E
NCBI Official Synonym Symbols	CBP; EIF4F; AUTS19; EIF4E1; eIF-4E; EIF4EL1
NCBI Protein Information	eukaryotic translation initiation factor 4E
UniProt Protein Name	Eukaryotic translation initiation factor 4E
UniProt Gene Name	EIF4E
UniProt Synonym Gene Names	EIF4EL1; EIF4F; eIF-4E; eIF4E

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/monoclonal/eif4e-recombinant-antibody-immunohistochemistry-western-blot-elisa/235538>