

MMP9 recombinant antibody

MMP9 Antibody



CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA235573	Rabbit	Human	Flow Cytometry, Functional Assay, Immunohistochemistry, ELISA

PRODUCT SYNONYMS

MMP9; Monoclonal Recombinant Antibody; MMP9 Antibody; Matrix metalloproteinase-9; MMP-9; 92 kDa gelatinase; 92 kDa type IV collagenase; Gelatinase B; GELB; 67 kDa matrix metalloproteinase-9; 82 kDa matrix metalloproteinase-9; CLG4B; MMP9 recombinant antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG
Purity/Purification	Affinity-Chromatography
Form/Format	Liquid; In phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Sequence Length	707
Applicable Applications for MMP9 recombinant antibody	FCM/FACS (Flow Cytometry), IHC (Immunohistochemistry), ELISA
Immunogen Species	Human
Immunogen	A synthesized peptide derived from human MMP9
Conjugate	Non-Conjugated
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.



IHC (Immunohistochemistry)
IHC image of CSB-RA014679A0HU diluted at 1:235 and staining in paraffin-embedded human breast cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

FCM/FACS (Flow Cytometry)
Overlay histogram showing Jurkat cells stained with CSB-RA014679A0HU (red line) at 1:50. The cells were fixed with 70% Ethylalcohol (18h) and then permeabilized with 0.3% Triton X-100 for 2 min. The cells were then incubated in 1x PBS /10% normal goat serum to block non-specific protein-protein interactions followed by primary antibody for 1 h at 4 degree C. The secondary antibody used was FITC goat anti-rabbit IgG (H+L) at 1/200 dilution for 1 h at 4 degree C. Control antibody (green line) was used under the same conditions. Acquisition of >10,000 events was performed.

ADDITIONAL INFORMATION

Related Product Information for MMP9 recombinant antibody	May play an essential role in local proteolysis of the extracellular matrix and in leukocyte migration. Could play a role in bone osteoclastic resorption. Cleaves KiSS1 at a Gly-[Leu bond. Cleaves type IV and type V collagen into large C-terminal three quarter fragments and shorter N-terminal one quarter fragments. Degrades fibronectin but not laminin or Pz-peptide.
Product Categories/Family for MMP9 recombinant antibody	Recombinant Antibody; Cardiovascular

NCBI AND UNIPROT INFORMATION

NCBI GI #	74272287
NCBI GeneID	4318
NCBI Accession #	NP_004985.2
NCBI GenBank Nucleotide #	NM_004994.3
UniProt Accession #	P14780; Q3LR70; Q8N725; Q9H4Z1; Q9UCJ9; Q9UCL1; Q9UDK2; B2R7V9
Molecular Weight	78,458 Da
NCBI Official Full Name	matrix metalloproteinase-9 preproprotein
NCBI Official Synonym Full Names	matrix metalloproteinase 9
NCBI Official Symbol	MMP9
NCBI Official Synonym Symbols	GELB; CLG4B; MMP-9; MANDP2
NCBI Protein Information	matrix metalloproteinase-9
UniProt Protein Name	Matrix metalloproteinase-9
UniProt Gene Name	MMP9
UniProt Synonym Gene Names	CLG4B; GELB

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/monoclonal/mmp9-recombinant-antibody-flow-cytometry-immunohistochemistry-elisa/235573>