

RAF1 recombinant antibody

Phospho-RAF1 (S259) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA235584	Rabbit	Human	Immunofluorescence, Immunohistochemistry, Western Blot, ELISA


PRODUCT SYNONYMS

RAF1; Monoclonal Recombinant Antibody; Phospho-RAF1 (S259) Antibody; RAF proto-oncogene serine/threonine-protein kinase; Proto-oncogene c-RAF; cRaf; Raf-1; RAF; RAF1 recombinant antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG
Purity/Purification	Affinity-Chromatography
Form/Format	Liquid; In phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Sequence Length	648
Applicable Applications for RAF1 recombinant antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen Species	Human
Immunogen	A synthesized peptide derived from human Phospho-RAF1 (S259)
Conjugate	Non-Conjugated
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IF (Immunofluorescence)

Immunofluorescence staining of HepG2 cells with CSB-RA019284A259pHU at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L).



IHC (Immunohistochemistry)

IHC image of CSB-RA019284A259pHU diluted at 1:100 and staining in paraffin-embedded human endometrial cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



WB (Western Blot)
 Western BlotPositive WB detected in?Hela whole cell lysate(treated with Calyculin A or EGF)All lanes?Phospho-RAF1 antibody at 1.18ug/mlSecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 73 KDaObserved band size: 73 KDa

ADDITIONAL INFORMATION

Related Product Information for RAF1 recombinant antibody	Serine/threonine-protein kinase that acts as a regulatory link between the membrane-associated Ras GTPases and the MAPK/ERK cascade, and this critical regulatory link functions as a switch determining cell fate decisions including proliferation, differentiation, apoptosis, survival and oncogenic transformation. RAF1 activation initiates a mitogen-activated protein kinase (MAPK) cascade that comprises a sequential phosphorylation of the dual-specific MAPK kinases (MAP2K1/MEK1 and MAP2K2/MEK2) and the extracellular signal-regulated kinases (MAPK3/ERK1 and MAPK1/ERK2). The phosphorylated form of RAF1 (on residues Ser-338 and Ser-339, by PAK1) phosphorylates BAD/Bcl2-antagonist of cell death at 'Ser-75'. Phosphorylates adenylyl cyclases: ADCY2, ADCY5 and ADCY6, resulting in their activation. Phosphorylates PPP1R12A resulting in inhibition of the phosphatase activity. Phosphorylates TNNT2/cardiac muscle troponin T. Can promote NF-kB activation and inhibit signal transducers involved in motility (ROCK2), apoptosis (MAP3K5/ASK1 and STK3/MST2), proliferation and angiogenesis (RB1). Can protect cells from apoptosis also by translocating to the mitochondria where it binds BCL2 and displaces BAD/Bcl2-antagonist of cell death. Regulates Rho signaling and migration, and is required for normal wound healing. Plays a role in the oncogenic transformation of epithelial cells via repression of the TJ protein, occludin (OCLN) by inducing the up-regulation of a transcriptional repressor SNAI2/SLUG, which induces down-regulation of OCLN. Restricts caspase activation in response to selected stimuli, notably Fas stimulation, pathogen-mediated macrophage apoptosis, and erythroid differentiation.
Product Categories/Family for RAF1 recombinant antibody	Recombinant Antibody ; Signal Transduction

NCBI AND UNIPROT INFORMATION

NCBI GI #	4506401
NCBI GeneID	5894
NCBI Accession #	NP_002871.1
NCBI GenBank Nucleotide #	NM_002880.3
UniProt Accession #	P04049
NCBI Official Full Name	RAF proto-oncogene serine/threonine-protein kinase isoform b
NCBI Official Synonym Full Names	Raf-1 proto-oncogene, serine/threonine kinase
NCBI Official Symbol	RAF1
NCBI Official Synonym Symbols	NS5; CRAF; Raf-1; c-Raf; CMD1NN
NCBI Protein Information	RAF proto-oncogene serine/threonine-protein kinase

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

