

Rabbit anti-Human HTRA1 Polyclonal Antibody | anti-HTRA1 antibody

HTRA1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA243546	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

HTRA1; Polyclonal Antibody; HTRA1 Antibody; Serine protease HTRA1 (EC 3.4.21.-) (High-temperature requirement A serine peptidase 1) (L56) (Serine protease 11); HTRA PRSS11; anti-HTRA1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G Purified
Form/Format	Liquid Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Applicable Applications for anti-HTRA1 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant Human Serine protease HTRA1 protein (23-248AA)
Immunogen Species	Human
Conjugate	Non-conjugated
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IF (Immunofluorescence)

Immunofluorescence staining of HepG2 cells with AAA243546 at 1:166, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IHC (Immunohistochemistry)

IHC image of AAA243546 diluted at 1:500 and staining in paraffin-embedded human placenta tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)

IHC image of AAA243546 diluted at 1:500 and staining in paraffin-embedded human kidney tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

ADDITIONAL INFORMATION

Product Categories/Family for anti-HTRA1 antibody	Signal Transduction
---	---------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	18202620
NCBI GeneID	5654
UniProt Accession #	Q92743; Q9UNS5; D3DRE4
NCBI Official Full Name	Serine protease HTRA1
NCBI Official Synonym Full Names	HtrA serine peptidase 1
NCBI Official Symbol	HTRA1
NCBI Official Synonym Symbols	L56; HtrA; ARMD7; ORF480; PRSS11; CARASIL
NCBI Protein Information	serine protease HTRA1; IGFBP5-protease; protease, serine, 11 (IGF binding); high-temperature requirement A serine peptidase 1
UniProt Protein Name	Serine protease HTRA1
UniProt Gene Name	HTRA1
UniProt Synonym Gene Names	HTRA; PRSS11
UniProt Entry Name	HTRA1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://www.aaabiotech.com/polyclonal/htra1-antibody-immunofluorescence-immunohistochemistry-elisa/243546>