

Mouse anti-Human CEACAM5 Monoclonal Antibody | anti-CEACAM5 antibody

CEACAM5 Monoclonal Antibody

CATALOG # AAA243702	HOST Mouse	REACTIVITY Human	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	----------------------	----------------------------	--


PRODUCT SYNONYMS

CEACAM5; Monoclonal Antibody; CEACAM5 Monoclonal Antibody; Carcinoembryonic antigen-related cell adhesion molecule 5 (Carcinoembryonic antigen) (CEA) (Meconium antigen 100) (CD antigen CD66e); CEA; anti-CEACAM5 antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG2b, Kappa
Purity/Purification	Affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.
Form/Format	Liquid; In Phosphate Buffered Saline, 50% Glycerol, 0.5% BSA and 0.02% Sodium Azide.
Applicable Applications for anti-CEACAM5 antibody	IHC (Immunohistochemistry), ELISA
Immunogen	Synthesized peptide derived from human carcinoembryonic antigen(CEA)
Immunogen Species	Human
Conjugate	Non-conjugated
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IHC (Immunohistochemistry)

IHC image of AAA243702 diluted at 1:100 and staining in paraffin-embedded human gastric cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-mouse IgG polymer labeled by HRP and visualized using 0.05% DAB.



IHC (Immunohistochemistry)

IHC image of AAA243702 diluted at 1:100 and staining in paraffin-embedded human colon cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-mouse IgG polymer labeled by HRP and visualized using 0.05% DAB.



IHC (Immunohistochemistry)

IHC image of AAA243702 diluted at 1:100 and staining in paraffin-embedded human tonsil tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-mouse IgG polymer labeled by HRP and visualized using 0.05% DAB.

ADDITIONAL INFORMATION

Product Categories/Family for anti-CEACAM5 antibody	Tags & Cell Markers
---	---------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	317373456
NCBI GeneID	1048
UniProt Accession #	P06731; H9KVA7
Molecular Weight	76,795 Da
NCBI Official Full Name	Carcinoembryonic antigen-related cell adhesion molecule 5
NCBI Official Synonym Full Names	carcinoembryonic antigen-related cell adhesion molecule 5
NCBI Official Symbol	CEACAM5
NCBI Official Synonym Symbols	CEA; CD66e
NCBI Protein Information	carcinoembryonic antigen-related cell adhesion molecule 5; meconium antigen 100
UniProt Protein Name	Carcinoembryonic antigen-related cell adhesion molecule 5
UniProt Gene Name	CEACAM5
UniProt Synonym Gene Names	CEA; CEA
UniProt Entry Name	CEAM5_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://www.aaabiotech.com/monoclonal/ceacam5-antibody-immunohistochemistry-elisa/243702>