

Rabbit anti-Human ATF4 Polyclonal Antibody | anti-ATF4 antibody

ATF4 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA243759	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

ATF4; Polyclonal Antibody; ATF4 Antibody; Cyclic AMP-dependent transcription factor ATF-4 (cAMP-dependent transcription factor ATF-4) (Activating transcription factor 4) (Cyclic AMP-responsive element-binding protein 2) (CREB-2) (cAMP-responsive element-binding protein 2) (DNA-binding protein TAXREB67) (Tax-responsive enhancer element-binding protein 67) (TaxREB67); CREB2 TXREB; anti-ATF4 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH7.4
Applicable Applications for anti-ATF4 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Conjugation	Non-conjugated
Immunogen	Recombinant Human Cyclic AMP-dependent transcription factor ATF-4 protein (20-200aa)
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IF (Immunofluorescence)

Immunofluorescence staining of HeLa cells at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IHC (Immunohistochemistry)

IHC image diluted at 1:100 and staining in paraffin-embedded human breast cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.05% DAB.

ADDITIONAL INFORMATION

Related Product Information for anti-ATF4 antibody	Transcriptional activator. Binds the cAMP response element (CRE) (consensus: 5'-GTGACGT[AC][AG]-3'), a sequence present in many viral and cellular promoters. Cooperates with FOXO1 in osteoblasts to regulate glucose homeostasis through suppression of beta-cell production and decrease in insulin production (By similarity). It binds to a Tax-responsive enhancer element in the long terminal repeat of HTLV-I. Regulates the induction of DDIT3/CHOP and asparagine synthetase (ASNS) in response to endoplasmic reticulum (ER) stress. In concert with DDIT3/CHOP, activates the transcription of TRIB3 and promotes ER stress-induced neuronal apoptosis by regulating the transcriptional induction of BBC3/PUMA. Activates transcription of SIRT4. Regulates the circadian expression of the core clock component PER2 and the serotonin transporter SLC6A4. Binds in a circadian time-dependent manner to the cAMP response elements (CRE) in the SLC6A4 and PER2 promoters and periodically activates the transcription of these genes. During ER stress response, activates the transcription of NLRP1, possibly in concert with other factors (PubMed:26086088).
Product Categories/Family for anti-ATF4 antibody	Signal Transduction

NCBI AND UNIPROT INFORMATION

NCBI GI #	33469976
NCBI GeneID	468
NCBI Accession #	NP_001666.2
NCBI GenBank Nucleotide #	NM_001675.2
UniProt Accession #	P18848; Q9UHQ1
Molecular Weight	40 kDa
NCBI Official Full Name	cyclic AMP-dependent transcription factor ATF-4
NCBI Official Synonym Full Names	activating transcription factor 4
NCBI Official Symbol	ATF4
NCBI Official Synonym Symbols	CREB2; TXREB; CREB-2; TAXREB67
NCBI Protein Information	cyclic AMP-dependent transcription factor ATF-4; DNA-binding protein TAXREB67; tax-responsive enhancer element B67; cAMP response element-binding protein 2; cAMP-dependent transcription factor ATF-4; cAMP-responsive element-binding protein 2; cyclic AMP-r
UniProt Protein Name	Cyclic AMP-dependent transcription factor ATF-4
UniProt Gene Name	ATF4
UniProt Synonym Gene Names	CREB2; TXREB; cAMP-dependent transcription factor ATF-4; CREB-2; cAMP-responsive element-binding protein 2; TaxREB67
UniProt Entry Name	ATF4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/atf4-antibody-immunofluorescence-immunohistochemistry-elisa/243759>