

Rabbit anti-Human HIST1H3A Polyclonal Antibody | anti-HIST1H3A antibody

Rabbit anti-Human HIST1H3A Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA244081	Rabbit	Human	Immunohistochemistry, ELISA


PRODUCT SYNONYMS

HIST1H3A; Polyclonal Antibody; Rabbit anti-Human HIST1H3A Polyclonal Antibody; Histone H3.1 (Histone H3/a) (Histone H3/b) (Histone H3/c) (Histone H3/d) (Histone H3/f) (Histone H3/h) (Histone H3/i) (Histone H3/j) (Histone H3/k) (Histone H3/l); HIST1H3B; HIST1H3C; HIST1H3D; HIST1H3E; HIST1H3F; HIST1H3G; HIST1H3H; HIST1H3I; HIST1H3J; H3FA; H3FL; H3FC; H3FB; H3FD; H3FI; H3FH; H3FK; H3FF; H3FJ; HIST1H3A (Ab-122) Antibody; anti-HIST1H3A antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid. Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH7.4
Applicable Applications for anti-HIST1H3A antibody	IHC (Immunohistochemistry), ELISA
Immunogen Species	Human
Immunogen	Synthesized peptide derived from Human Histone H3.1 protein (120-132aa)
Conjugation	Non-conjugated
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)

IHC image of diluted at 1:20 and staining in paraffin-embedded human liver tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)

IHC image of diluted at 1:20 and staining in paraffin-embedded human small intestine tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

ADDITIONAL INFORMATION

Related Product Information for anti-HIST1H3A antibody	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
Product Categories/Family for anti-HIST1H3A antibody	Epigenetics and Nuclear Signaling

NCBI AND UNIPROT INFORMATION

NCBI GI #	4504281
NCBI GeneID	8350
NCBI Accession #	NP_003520.1
NCBI GenBank Nucleotide #	NM_003529.2
UniProt Accession #	P68431; P02295; P02296; P16106; Q6ISV8; Q6NWP8; Q6NWP9; Q6NXU4; Q71DJ3; Q93081; A0PJT7; A5PLR1
Molecular Weight	15,404 Da
NCBI Official Full Name	histone H3.1
NCBI Official Synonym Full Names	histone cluster 1, H3a
NCBI Official Symbol	HIST1H3A
NCBI Official Synonym Symbols	H3/A; H3FA
NCBI Protein Information	histone H3.1; histone H3/a; histone 1, H3a; H3 histone family, member A
UniProt Protein Name	Histone H3.1
UniProt Gene Name	HIST1H3A
UniProt Synonym Gene Names	H3FA
UniProt Entry Name	H31_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/hist1h3a-antibody-immunohistochemistry-elisa/244081>