Rabbit anti-Human HIST1H4A Polyclonal Antibody | anti-HIST1H4A antibody

Rabbit anti-Human HIST1H4A Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA244089	**Rabbit**	**Human**	**Immunohistochemistry, ELISA**

PRODUCT SYNONYMS

HIST1H4A; Polyclonal Antibody; Rabbit anti-Human HIST1H4A Polyclonal Antibody; Histone H4; HIST1H4B; HIST1H4C; HIST1H4D; HIST1H4E; HIST1H4F; HIST1H4H; HIST1H4I; HIST1H4J; HIST1H4K; HIST1H4L; HIST2H4A; HIST2H4B; HIST4H4; H4/A H4FA; H4/I H4FI; H4/G H4FG; H4/B H4FB; H4/J H4FJ; H4/C H4FC; H4/H H4FH; H4/M H4FM; H4/E H4FE; H4/D H4FD; H4/K H4FK; H4/N H4F2 H4FN HIST2H4; H4/O H4FO; HIST1H4A (Ab-77) Antibody; anti-HIST1H4A antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid. Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH7.4
Applicable Applications for anti-HIST1H4A antibody	IHC (Immunohistochemistry), ELISA
Immunogen Species	Human
Immunogen	Synthesized peptide derived from Human Histone H4 protein (74-85aa)
Conjugation	Non-conjugated
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohiostchemistry)
IHC image of diluted at 1:20 and staining in paraffin-embedded human liver cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)
IHC image of diluted at 1:20 and staining in paraffin-embedded human colon cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

ADDITIONAL INFORMATION

Related Product Information for anti-HIST1H4A antibody	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
Product Categories/Family for anti-HIST1H4A antibody	Epigenetics and Nuclear Signaling

NCBI AND UNIPROT INFORMATION

NCBI GI #	77539758
NCBI GeneID	554313
NCBI Accession #	NP_001029249.1
NCBI GenBank Nucleotide #	NM_001034077.4
UniProt Accession #	P62805; P02304; P02305; Q6DRA9; Q6FGB8; Q6NWP7; A2VCL0
Molecular Weight	11,367 Da
NCBI Official Full Name	histone H4
NCBI Official Synonym Full Names	histone cluster 2, H4b
NCBI Official Symbol	HIST2H4B
NCBI Official Synonym Symbols	H4/o
NCBI Protein Information	histone H4; histone 2, H4b
UniProt Protein Name	Histone H4
UniProt Gene Name	**HIST1H4A**
UniProt Synonym Gene Names	H4/A; H4FA
UniProt Entry Name	H4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: https://www.aaabiotech.com/polyclonal/hist1h4a-antibody-immunohistochemistry-elisa/244089