

Rabbit NSE/ENO2 Polyclonal Antibody | anti-ENO2 antibody

Anti-NSE/ENO2 Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA254171	Rabbit	Human Predicted to React with: Rat (Species predicted to react based on 100% sequence homology)	Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

NSE/ENO2; Polyclonal Antibody; Anti-NSE/ENO2 Antibody; Rabbit Polyclonal; NSE(Gamma-enolase) Antibody; Rabbit PAb; Antigen Affinity Purified; enolase 2 (gamma; neuronal); Anti-HEL-S-279 Antibody; Anti-NSE Antibody; anti-ENO2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human Predicted to React with: Rat (Species predicted to react based on 100% sequence homology)
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human NSE/ENO2
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; 0.2um filtered solution in PBS
Applicable Applications for anti-ENO2 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthetic peptide corresponding to the N-terminus of the human NSE(Gamma-enolase)
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with A synthetic peptide corresponding to the N-terminus of the human NSE(Gamma-enolase), and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Preservative-Free. Avoid repeated freeze-thaw cycles.

IMAGES




WB (Western Blot)

Anti-NSE(Gamma-enolase) rabbit polyclonal antibody at 1:500 dilutionLane A: HepG2 Whole Cell LysateLysates/proteins at 30 ug per lane.SecondaryGoat Anti-Rabbit IgG H&L (Dylight800) at 1/10000 dilution.Developed using the Odyssey technique.Performed under reducing conditions.Predicted band size:47 kDaObserved band size:55 kDa



IHC (Immunohistochemistry)

Immunochemical staining of human NSE in human brain with rabbit polyclonal antibody (1:10000, formalin-fixed paraffin embedded sections).



IHC (Immunohistochemistry)

Immunochemical staining of human NSE in human pancreas with rabbit polyclonal antibody (1:10000, formalin-fixed paraffin embedded sections).

ADDITIONAL INFORMATION

Related Product Information for anti-ENO2 antibody	The combination of silencing ENO2 and 2-deoxyglucose (2-DG) synergistically inhibited leukemia cell survival. ENO2 may be a biological marker for monitoring chemotherapeutic efficacy and relapse in ALL. Reduced ENO2 expression may be a biomarker for a subset of autistic children. Neuron specific enolase (ENO2, gamma-enolase) has been used as a biomarker to help identify neuroendocrine differentiation in breast cancer.
References	Pechumer H, et al. (1994) Detection of neuron-specific gamma-enolase messenger ribonucleic acid in normal human leukocytes by polymerase chain reaction amplification with nested primers. Lab Invest. 69 (6): 743-9.Craig SP, et al. (1991) Localisation of neurone-specific enolase (ENO2) to 12p13. Cytogenet Cell Genet. 54 (1-2): 71-3.Muley T, et al. (2003) Technical performance and diagnostic utility of the new Elecsys neuron-specific enolase enzyme immunoassay. Clin Chem Lab Med. 41 (1): 95-103.

NCBI AND UNIPROT INFORMATION

NCBI GI #	5803011
NCBI GeneID	2026
NCBI Accession #	NP_001966.1
NCBI GenBank Nucleotide #	NM_001975.2
UniProt Accession #	P09104; Q96J33
Molecular Weight	47,269 Da

NCBI Official Full Name	gamma-enolase
NCBI Official Synonym Full Names	enolase 2 (gamma, neuronal)
NCBI Official Symbol	ENO2
NCBI Official Synonym Symbols	NSE
NCBI Protein Information	gamma-enolase; neural enolase; neuron-specific enolase; neurone-specific enolase; neuron specific gamma enolase; 2-phospho-D-glycerate hydrolyase; 2-phospho-D-glycerate hydro-lyase
UniProt Protein Name	Gamma-enolase
UniProt Gene Name	ENO2
UniProt Synonym Gene Names	NSE
UniProt Entry Name	ENOG_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/nse-eno2-antibody-immunohistochemistry-western-blot-elisa/254171>