

Rabbit anti-Human CHEK2 Polyclonal Antibody | anti-CHEK2 antibody

Anti-CHEK2 Antibody, Rabbit Polyclonal



CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA254381	Rabbit	Human	Immunoprecipitation, Western Blot

PRODUCT SYNONYMS

CHEK2; Polyclonal Antibody; Anti-CHEK2 Antibody; Rabbit Polyclonal; CHEK2 Antibody; Rabbit PAb; Antigen Affinity Purified; checkpoint kinase 2; Anti-CDS1 Antibody; Anti-CHK2 Antibody; Anti-hCds1 Antibody; Anti-HuCds1 Antibody; Anti-LFS2 Antibody; Anti-PP1425 Antibody; Anti-RAD53 Antibody; anti-CHEK2 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human CHEK2
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-CHEK2 antibody	IP (Immunoprecipitation), WB (Western Blot)
Immunogen	A synthetic peptide corresponding to the N-terminus of the Human CHEK2
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with a synthetic peptide corresponding to the N-terminus of the Human CHEK2, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.



IP (Immunoprecipitation)
CHEK2 was immunoprecipitated using: Lane A: 0.5 mg Jurkat Whole Cell Lysate 4 uL anti-CHEK2 rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G. Primary antibody: Anti-CHEK2 rabbit polyclonal antibody, at 1:100 dilution Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 61 kDa Observed band size: 61 kDa

WB (Western Blot)
Anti-CHEK2 rabbit polyclonal antibody at 1:500 dilution Lane A: Jurkat Whole Cell Lysate Lysates/proteins at 30 ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 61 kDa Observed band size: 61 kDa

ADDITIONAL INFORMATION

Related Product Information for anti-CHEK2 antibody	In response to DNA damage and replication blocks, cell cycle progression is halted through the control of critical cell cycle regulators. The protein encoded by CHEK2 gene is a cell cycle checkpoint regulator and putative tumor suppressor. It contains a forkhead-associated protein interaction domain essential for activation in response to DNA damage and is rapidly phosphorylated in response to replication blocks and DNA damage. When activated, the encoded CHEK2 protein is known to inhibit CDC25C phosphatase, preventing entry into mitosis, and has been shown to stabilize the tumor suppressor protein p53, leading to cell cycle arrest in G1. In addition, this protein interacts with and phosphorylates BRCA1, allowing BRCA1 to restore survival after DNA damage. Mutations in this gene have been linked with Li-Fraumeni syndrome, a highly penetrant familial cancer phenotype usually associated with inherited mutations in TP53. Also, mutations in CHEK2s gene are thought to confer a predisposition to sarcomas, breast cancer, and brain tumors. This nuclear protein is a member of the CDS1 subfamily of serine/threonine protein kinases. Several transcript variants encoding different isoforms have been found for this gene.
Product Categories/Family for anti-CHEK2 antibody	Cancer Drug Targets Tumor Suppressors; p53 Pathway
References	Bogdanova N, et al. (2005) Association of two mutations in the CHEK2 gene with breast cancer. Cancer Genetics. 116(2) : 263-6. Dong XY, et al. (2003) Mutations in CHEK2 associated with prostate cancer risk. The American journal of human genetics. 72(2) 270-80. Massink MPG, Kooi IE, Martens JWM, Waisfisz Q, Meijers-Heijboer H. Genomic profiling of CHEK2*1100delC-mutated breast carcinomas. BMC Cancer. 2015;15:877. doi:10.1186/s12885-015-1880-y.

NCBI AND UNIPROT INFORMATION

NCBI GI #	54112407
NCBI GeneID	11200
NCBI Accession #	NP_001005735.1
NCBI GenBank Nucleotide #	NM_001005735.1
UniProt Accession #	O96017; Q6QA03; Q6QA04; Q6QA05; Q6QA06; Q6QA07; Q6QA08; Q6QA10; A8K3Y9; B7ZBF3; B7ZBF4; B7ZBF5
Molecular Weight	60,915 Da
NCBI Official Full Name	serine/threonine-protein kinase Chk2 isoform c
NCBI Official Synonym Full Names	checkpoint kinase 2

NCBI Official Symbol	CHEK2
NCBI Official Synonym Symbols	CDS1; CHK2; LFS2; RAD53; hCds1; HuCds1; PP1425
NCBI Protein Information	serine/threonine-protein kinase Chk2; cds1 homolog; CHK2 checkpoint homolog; checkpoint-like protein CHK2
UniProt Protein Name	Serine/threonine-protein kinase Chk2
UniProt Gene Name	CHEK2
UniProt Synonym Gene Names	CDS1; CHK2; RAD53; Hucds1; hCds1
UniProt Entry Name	CHK2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/chek2-antibody-immunoprecipitation-western-blot-elisa/254381>