

Mouse anti-Human CD23 Monoclonal Antibody | anti-FCER2 antibody

Anti-CD23 Antibody, Mouse Monoclonal



CATALOG # AAA254769	HOST Mouse	REACTIVITY Human	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	----------------------	----------------------------	--

PRODUCT SYNONYMS

CD23; Monoclonal Antibody; Anti-CD23 Antibody; Mouse Monoclonal; CD23/FCER2 Antibody; Mouse MAb; Fc fragment of IgE receptor II; Anti-BLAST-2 Antibody; Anti-CD23A Antibody; Anti-CLEC-6 Antibody; Anti-CLEC4J Antibody; Anti-CLEC6 Antibody; Anti-CLECSF8 Antibody; Anti-FCE2 Antibody; Anti-IGEBF Antibody; Anti-MCL Antibody; Anti-MGC40078 Antibody; Anti-MPCL Antibody; anti-FCER2 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse IgG2a
Clone Number	05
Specificity	Human CD23/FCER2. No cross-reactivity in ELISA with Human cell lysate (293 cell line)
Purity/Purification	Protein A
Form/Format	Liquid; 0.2um filtered solution in PBS
Applicable Applications for anti-FCER2 antibody	IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant Human CD23/FCER2 protein
Conjugation	Unconjugated
Preparation	This antibody was produced from a hybridoma resulting from the fusion of a mouse myeloma with B cells obtained from a mouse immunized with purified, recombinant Human CD23/FCER2 (rh CD23/FCER2). The IgG fraction of the cell culture supernatant was purified by Protein A affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Preservative-Free. Avoid repeated freeze-thaw cycles.



IHC (Immunohistochemistry)
Immunochemical staining of human CD23 in human spleen with mouse monoclonal antibody (1:60, formalin-fixed paraffin embedded sections).

IHC (Immunohistochemistry)
Immunochemical staining of human CD23 in human tonsil with mouse monoclonal antibody (1:60, formalin-fixed paraffin embedded sections).

ADDITIONAL INFORMATION

Related Product Information for anti-FCER2 antibody	Fc fragment of IgE, low affinity II, receptor for (CD23) or CD23 antigen is a member of the cluster of differentiation family. The cluster of differentiation (cluster of designation) (often abbreviated as CD) is a protocol used for the identification and investigation of cell surface molecules present on white blood cells initially but found in almost any kind of cell of the body, providing targets for immunophenotyping of cells. Physiologically, CD molecules can act in numerous ways, often acting as receptors or ligands (the molecule that activates a receptor) important to the cell. A signal cascade is usually initiated, altering the behavior of the cell (see cell signaling). Some CD proteins do not play a role in cell signaling, but have other functions, such as cell adhesion. CD23/FCER2 is a B-cell specific antigen, and a low-affinity receptor for IgE. It has essential roles in B cell growth and differentiation, and the regulation of IgE production. This protein also exists as a soluble secreted form, then functioning as a potent mitogenic growth factor. Increased levels of soluble CD23/FCER2 cause the recruitment of non-sensitised B-cells in the presentation of antigen peptides to allergen-specific B-cells, therefore increasing the production of allergen specific IgE. IgE, in turn, is known to upregulate the cellular expression of CD23 and Fc epsilon RI (high-affinity IgE receptor).
Product Categories/Family for anti-FCER2 antibody	Fc Receptors ITIM/ITAM Immunoreceptors and Related Molecules
References	Aubry JP, et al. (1992) CD21 is a ligand for CD23 and regulates IgE production. Nature. 358(6386): 505-7.Punnonen J, et al. (1993) Interleukin 13 induces interleukin 4-independent IgG4 and IgE synthesis and CD23 expression by human B cells. Proc Natl Acad Sci U S A. 90(8): 3730-4.Yu P, et al. (1994) Negative feedback regulation of IgE synthesis by murine CD23. Nature. 369(6483): 753-6.

NCBI AND UNIPROT INFORMATION

NCBI GI #	119862
NCBI GeneID	2208
UniProt Accession #	P06734
Molecular Weight	321
NCBI Official Full Name	Low affinity immunoglobulin epsilon Fc receptor
NCBI Official Synonym Full Names	Fc fragment of IgE, low affinity II, receptor for (CD23)
NCBI Official Symbol	FCER2
NCBI Official Synonym Symbols	CD23; FCE2; CD23A; IGEBF; CLEC4J; BLAST-2
NCBI Protein Information	low affinity immunoglobulin epsilon Fc receptor; CD23 antigen; fc-epsilon-RII; lymphocyte IgE receptor; immunoglobulin E-binding factor; C-type lectin domain family 4, member J
UniProt Protein Name	Low affinity immunoglobulin epsilon Fc receptor
UniProt Gene Name	FCER2

UniProt Synonym Gene Names	CD23A; CLEC4J; FCE2; IGEBF
UniProt Entry Name	FCER2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/monoclonal/cd23-antibody-clone-05-immunohistochemistry-elisa/254769>