

Rabbit anti-Human MUC1 Polyclonal Antibody | anti-MUC1 antibody

Anti-MUC1 Antibody, Rabbit Polyclonal


CATALOG # AAA255523	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	-----------------------	----------------------------	--

PRODUCT SYNONYMS


MUC1; Polyclonal Antibody; Anti-MUC1 Antibody; Rabbit Polyclonal; MUC-1/Mucin 1/CD227 Antibody; Rabbit PAb; Antigen Affinity Purified; mucin 1; cell surface associated; Anti-ADMCKD Antibody; Anti-ADMCKD1 Antibody; Anti-CA15-3 Antibody; Anti-CD227 Antibody; Anti-EMA Antibody; Anti-H23AG Antibody; Anti-KL-6 Antibody; Anti-MAM6 Antibody; Anti-MCD Antibody; Anti-MCKD Antibody; Anti-MCKD1 Antibody; Anti-MUC-1 Antibody; Anti-MUC-1/SEC Antibody; Anti-MUC-1/X Antibody; Anti-MUC1/ZD Antibody; Anti-Mucin 1 Antibody; Anti-PEM Antibody; Anti-PEMT Antibody; Anti-PUM Antibody; anti-MUC1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human MUC-1/Mucin 1/CD227
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; 0.2um filtered solution in PBS
Applicable Applications for anti-MUC1 antibody	IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant Human MUC-1/Mucin 1/CD227 protein
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with purified, recombinant Human MUC-1/Mucin 1/CD227 (rh MUC-1/Mucin 1/CD227;NP_001018016.1; Met1-Gly167). MUC-1/Mucin 1/CD227 specific IgG was purified by Human MUC-1/Mucin 1/CD227 affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Preservative-Free. Avoid repeated freeze-thaw cycles.



IHC (Immunohistochemistry)
Immunochemical staining of human MUC1 in human breast carcinoma with rabbit polyclonal antibody (1:5000, formalin-fixed paraffin embedded sections).



IHC (Immunohistochemistry)
Immunochemical staining of human MUC1 in human ovarian cancer with rabbit polyclonal antibody (1:5000, formalin-fixed paraffin embedded sections).

ADDITIONAL INFORMATION

Related Product Information for anti-MUC1 antibody	Mucin 1, cell surface-associated (MUC1) or polymorphic epithelial mucin (PEM) is a membrane-bound protein that is a member of the mucin family. Mucins are O-glycosylated proteins that play an essential role in forming protective mucous barriers on epithelial surfaces. These proteins also play a role in intracellular signaling. This protein is expressed on the apical surface of epithelial cells that line the mucosal surfaces of many different tissues including lung, breast stomach, and pancreas. MUC-1/CC1/CD227 Exclusively located in the apical domain of the plasma membrane of highly polarized epithelial cells. After endocytosis, internalized, and recycled to the cell membrane. This protein is proteolytically cleaved into alpha and beta subunits that form a heterodimeric complex. The N-terminal alpha subunit functions in cell-adhesion and the C-terminal beta subunit is involved in cell signaling. Overexpression, aberrant intracellular localization, and changes in glycosylation of this protein have been associated with carcinomas. The alpha subunit has cell adhesive properties. MUC-1/CC1/CD227 Can act both as an adhesion and an anti-adhesion protein. This protein May provide a protective layer on epithelial cells against bacterial and enzyme attack. The beta subunit contains a C-terminal domain which is involved in cell signaling, through phosphorylations and protein-protein interactions. MUC-1/CC1/CD227 participated in modulates signaling in ERK, SRC, and NF-kappa-B pathways. In the activated T-cells, MUC-1/CC1/CD227 influences directly or indirectly the Ras/MAPK pathway. MUC-1/CC1/CD227 Promotes tumor progression and regulates TP53-mediated transcription and determines cell fate in the genotoxic stress response. Binds, together with KLF4, the PE21 promoter element of TP53 and represses TP53 activity.
Product Categories/Family for anti-MUC1 antibody	Cancer Drug Targets; EGFR Signaling Pathway
References	Brayman M, et al. (2004) MUC1: a multifunctional cell surface component of reproductive tissue epithelia. Reprod Biol Endocrinol 2: 4. Schroeder J A, et al. (2001) Transgenic MUC1 interacts with epidermal growth factor receptor and correlates with mitogen-activated protein kinase activation in the mouse mammary gland. J Biol Chem. 276 (16): 13057-64. Li Y, et al. (2001) The epidermal growth factor receptor regulates interaction of the human DF3/MUC1 carcinoma antigen with c-Src and beta-catenin. J Biol Chem. 276 (38): 35239-42.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.