

Rabbit anti-Human Dynamitin/DCTN2 Polyclonal Antibody | anti-DCTN2 antibody

Anti-Dynamitin/DCTN2 Antibody, Rabbit Polyclonal


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA255984	Rabbit	Human	Immunofluorescence, Immunocytochemistry, Immunohistochemistry

PRODUCT SYNONYMS

Dynamitin/DCTN2; Polyclonal Antibody; Anti-Dynamitin/DCTN2 Antibody; Rabbit Polyclonal; Dynamitin/DCTN2 Antibody; Rabbit PAb; Antigen Affinity Purified; dynactin 2 (p50); Anti-DCTN50 Antibody; Anti-DYNAMITIN Antibody; Anti-HEL-S-77 Antibody; Anti-RBP50 Antibody; anti-DCTN2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human Dynamitin/DCTN2
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-DCTN2 antibody	IF (Immunofluorescence), ICC (Immunocytochemistry), IHC (Immunohistochemistry)
Immunogen	E Coli-derived Human Dynamitin/DCTN2 fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human Dynamitin/DCTN2 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.



IF (Immunofluorescence)

Immunofluorescence staining of DCTN2 in A431 cells. Cells were fixed with 4% PFA, permeabilized with 0.1% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human DCTN2 polyclonal antibody (dilution ratio 1:200) at 4 degree C overnight. Then cells were stained with the Alexa Fluor488-conjugated Goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI (blue). Positive staining was localized to Cytoplasm.



IHC (Immunohistochemistry)

Immunochemical staining of human DCTN2 in human kidney with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/dynamitin-dctn2-antibody-immunofluorescence-immunocytochemistry-immunohistochemistry-elisa/255984>