

Rabbit anti-Human CDC23 Polyclonal Antibody | anti-CDC23 antibody

Anti-CDC23 Antibody, Rabbit Polyclonal


CATALOG # AAA256020	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunoprecipitation, Western Blot
-------------------------------	-----------------------	----------------------------	--

PRODUCT SYNONYMS

CDC23; Polyclonal Antibody; Anti-CDC23 Antibody; Rabbit Polyclonal; CDC23 Antibody; Rabbit PAb; Antigen Affinity Purified; cell division cycle 23; Anti-ANAPC8 Antibody; Anti-APC8 Antibody; Anti-CUT23 Antibody; anti-CDC23 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human CDC23
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-CDC23 antibody	IP (Immunoprecipitation), WB (Western Blot)
Immunogen	E Coli-derived Human CDC23 fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human CDC23 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.



IP (Immunoprecipitation)

CDC23 was immunoprecipitated using: Lane A: 0.5 mg HeLa Whole Cell Lysate, 4 μ L anti-CDC23 rabbit polyclonal antibody and 60 μ g of Immunomagnetic beads Protein A/G. Primary antibody: Anti-CDC23 rabbit polyclonal antibody, at 1:100 dilution. Secondary antibody: Clean-Blot IP Detection Reagent (HRP) at 1:1000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 68 kDa. Observed band size: 60 kDa.



WB (Western Blot)

Anti-CDC23 rabbit polyclonal antibody at 1:500 dilution. Lane A: HeLa Whole Cell Lysate. Lane B: NCI-H460 Whole Cell Lysate. Lysates/proteins at 30 μ g per lane. Secondary: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 68 kDa. Observed band size: 60 kDa. (We are unsure as to the identity of these extra bands.)

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.