

Rabbit anti-Human FBXO7 Polyclonal Antibody | anti-FBXO7 antibody

Anti-FBXO7 Antibody, Rabbit Polyclonal

CATALOG # AAA256372	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

FBXO7; Polyclonal Antibody; Anti-FBXO7 Antibody; Rabbit Polyclonal; FBXO7 Antibody; Rabbit PAb; Antigen Affinity Purified; F-box protein 7; Anti-FBX Antibody; Anti-FBXO7 Antibody; Anti-FBX7 Antibody; Anti-PARK15 Antibody; Anti-PKPS Antibody; anti-FBXO7 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human FBXO7
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-FBXO7 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	E Coli-derived Human FBXO7 fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human FBXO7 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunochemical staining of human FBXO7 in human testis with rabbit polyclonal antibody at 1:300 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunochemical staining of human FBXO7 in human large intestine with rabbit polyclonal antibody at 1:300 dilution, formalin-fixed paraffin embedded sections.



WB (Western Blot)

Anti-FBXO7 rabbit polyclonal antibody at 1:500 dilution
 Lane A: MCF7 Whole Cell Lysate
 Lane B: HL-60 Whole Cell Lysate
 Lane C: U-251 MG Whole Cell Lysate
 Lysates/proteins at 30 ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 59 kDa Observed band size: 59 kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/fbxo7-antibody-immunohistochemistry-western-blot-elisa/256372>