

Rabbit anti-Human TTF2 Polyclonal Antibody | anti-TTF2 antibody

Anti-TTF2 Antibody, Rabbit Polyclonal

CATALOG # AAA256396	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunoprecipitation, Western Blot
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

TTF2; Polyclonal Antibody; Anti-TTF2 Antibody; Rabbit Polyclonal; TTF2 Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-HuF2 Antibody; anti-TTF2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human TTF2
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-TTF2 antibody	IP (Immunoprecipitation), WB (Western Blot)
Immunogen	E Coli-derived Human TTF2 fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human TTF2 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES



IP (Immunoprecipitation)

TTF2 was immunoprecipitated using: Lane A: 0.5 mg Jurkat Whole Cell Lysate, 4 uL anti-TTF2 rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G. Primary antibody: Anti-TTF2 rabbit polyclonal antibody, at 1:100 dilution. Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 130 kDa. Observed band size: 150 kDa.



WB (Western Blot)

Anti-TTF2 rabbit polyclonal antibody at 1:500 dilution. Lane A: Jurkat Whole Cell Lysate. Lysates/proteins at 30 ug per lane. Secondary: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 130 kDa. Observed band size: 130 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-TTF2 antibody	Papillary thyroid cancer (PTC), one of the most common malignant thyroid tumors, exists widely in the thyroid of adolescents. Thyroid transcription factor 1 (TTF1) and 2 (TTF2) were thyroid-specific transcription factors, and regulated expression of the thyroid-specific genes. TTF2 is an RNA polymerase II termination factor that is responsible for mitotic repression of transcription elongation.
--	---

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/ttf2-antibody-immunoprecipitation-western-blot-elisa/256396>