

Rabbit anti-Human SS18L2 Polyclonal Antibody | anti-SS18L2 antibody

Anti-SS18L2 Antibody, Rabbit Polyclonal

CATALOG # AAA256433	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

SS18L2; Polyclonal Antibody; Anti-SS18L2 Antibody; Rabbit Polyclonal; SS18L2 Antibody; Rabbit PAb; Antigen Affinity Purified; synovial sarcoma translocation gene on chromosome 18-like 2; Anti-KIAA-iso Antibody; anti-SS18L2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human SS18L2
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-SS18L2 antibody	IHC (Immunohistochemistry)
Immunogen	E Coli-derived Human SS18L2 fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human SS18L2 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Immunochemical staining of human SS18L2 in human gallbladder with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunochemical staining of human SS18L2 in human stomach with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.

ADDITIONAL INFORMATION

Related Product Information for anti-SS18L2 antibody	SS18L2 (SS18 Like 2, also known as SYT homolog 2) is a Protein Coding gene. The SS18L2 gene, located on 3p22.1, is conserved in Rhesus monkey, dog, cow, mouse, rat, chicken, zebrafish, and frog. SS18L2 belongs to the SS18 family and is homologous to SS18. An important paralog of this gene is SS18L1. SS18L2 is widely expressed in the lymph node, prostate, and other tissues. The highly conserved synovial sarcoma-associated protein SS18 is thought to act as a transcriptional co-activator through interactions with various proteins involved in (epigenetic) gene regulation. The SS18 SNH domain appears to act as a major interface for these protein-protein interactions.
References	Pech MF, et al. (2019) Systematic identification of cancer cell vulnerabilities to natural killer cell-mediated immune surveillance. Elife 8: e47362.de Bruijn DR, et al. (2006) Common origin of the human synovial sarcoma associated ss18 and ss18l1 gene loci. Cytogenet Genome Res 112 (3-4): 222-226.de Bruijn DR, et al. (2001) Mapping and characterization of the mouse and human ss18 genes, two human ss18-like genes and a mouse ss18 pseudogene. Cytogenet Cell Genet 92 (3-4): 310-319.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/ss18l2-antibody-immunohistochemistry-elisa/256433>