

Rabbit anti-Human ACIN1 Polyclonal Antibody | anti-ACIN1 antibody

Anti-ACIN1 Antibody, Rabbit Polyclonal

CATALOG # AAA256737	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

ACIN1; Polyclonal Antibody; Anti-ACIN1 Antibody; Rabbit Polyclonal; ACIN1 Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-ACINUS Antibody; Anti-ACN Antibody; Anti-fSAP152 Antibody; anti-ACIN1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human ACIN1
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-ACIN1 antibody	IHC (Immunohistochemistry)
Immunogen	E Coli-derived Human ACIN1 fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human ACIN1 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Immunochemical staining of human ACIN1 in human placenta with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunochemical staining of human ACIN1 in human brain with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.

ADDITIONAL INFORMATION

Related Product Information for anti-ACIN1 antibody	Apoptotic chromatin condensation inducer in the nucleus (also referred as ACIN1) was first characterized as an RNA-binding protein involved in apoptosis. ACIN1 is an auxiliary component of the exon junction complex (EJC) which is assembled throughout pre-messenger RNA splicing. The human ACIN1 gene gives rise to three protein isoforms, termed Acinus-L, Acinus-S-, and Acinus-S, with all containing an RRM domain and a C-terminal RS-like domain.
References	Xue L, et al. (2018) [expression and significance of acin1 mrna in platelets of lung cancer]. Zhongguo Fei Ai Za Zhi 21 (9): 677-681.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/acin1-antibody-immunohistochemistry-elisa/256737>