

Rabbit anti-Human ZNF782 Polyclonal Antibody | anti-ZNF782 antibody

Anti-ZNF782 Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA256839	Rabbit	Human	Immunofluorescence, Immunocytochemistry, Immunohistochemistry


PRODUCT SYNONYMS

ZNF782; Polyclonal Antibody; Anti-ZNF782 Antibody; Rabbit Polyclonal; ZNF782 Antibody; Rabbit PAb; Antigen Affinity Purified; anti-ZNF782 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human ZNF782
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-ZNF782 antibody	IF (Immunofluorescence), ICC (Immunocytochemistry), IHC (Immunohistochemistry)
Immunogen	E Coli-derived Human ZNF782 fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human ZNF782 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescence staining of ZNF782 in A431 cells. Cells were fixed with 4% PFA, permeabilized with 0.1% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human ZNF782 polyclonal antibody (dilution ratio 1:200) at 4 degree C overnight. Then cells were stained with the Alexa Fluor488-conjugated Goat Anti-rabbit IgG secondary antibody (green). Positive staining was localized to Nucleus.



IHC (Immunohistochemistry)

Immunochemical staining of human ZNF782 in human brain with rabbit polyclonal antibody at 1:2000 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunochemical staining of human ZNF782 in human gallbladder with rabbit polyclonal antibody at 1:2000 dilution, formalin-fixed paraffin embedded sections.

ADDITIONAL INFORMATION

Related Product Information for anti-ZNF782 antibody	ZNF782 (Zinc Finger Protein 782) is located on human chromosome 9q22.33. The ZNF782 protein is ubiquitously expressed in testis, thyroid and other tissues. Structurally, the encoded protein is reported to be 80904 Da in mass. The ZNF782 gene is conserved in chimpanzee, dog, cow, and frog. 68 organisms have orthologs with human gene ZNF782. ZNF782 is related to nucleic acid binding and DNA-binding transcription factor activity. ZNF782 may be involved in transcriptional regulation. An important paralog of this gene is ZNF300.
References	Hamada H, et al. (2019) Characterization of the human zinc finger nfx-1-type containing 1 encoding znfx1 gene and its response to 12-o-tetradecanoyl-13-acetate in hl-60 cells. Int J Oncol 55 (4): 896-904.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/znf782-antibody-immunofluorescence-immunocytochemistry-immunohistochemistry-elisa/256839>