

Rabbit anti-Human BEND7 Polyclonal Antibody | anti-BEND7 antibody

Anti-BEND7 Antibody, Rabbit Polyclonal

CATALOG # AAA256856	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

BEND7; Polyclonal Antibody; Anti-BEND7 Antibody; Rabbit Polyclonal; BEND7 Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-C10orf30 Antibody; anti-BEND7 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human BEND7
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-BEND7 antibody	IHC (Immunohistochemistry)
Immunogen	E Coli-derived Human BEND7 fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human BEND7 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Immunochemical staining of human BEND7 in human liver with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunochemical staining of human BEND7 in human kidney with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.

ADDITIONAL INFORMATION

Related Product Information for anti-BEND7 antibody	The BEN domain is predicted to function as an adaptor for the higher-order structuring of chromatin, and recruitment of chromatin-modifying factors in transcriptional regulation. It has been suggested to mediate protein-DNA and protein-protein interactions during chromatin organisation and transcription. The presence of BEN domains in a poxviral early virosomal protein and in polydnal proteins also suggests a possible role in the organisation of viral DNA during replication or transcription. They are generally linked to other globular domains with functions related to transcriptional regulation and chromatin structure, such as BTB, C4DM, and C2H2 fingers.
References	Dai Q, et al. (2013) The ben domain is a novel sequence-specific DNA-binding domain conserved in neural transcriptional repressors. Genes Dev 27 (6): 602-614.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/bend7-antibody-immunohistochemistry-elisa/256856>