

Rabbit anti-Human ZNF101 Polyclonal Antibody | anti-ZNF101 antibody

Anti-ZNF101 Antibody, Rabbit Polyclonal

CATALOG # AAA256983	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

ZNF101; Polyclonal Antibody; Anti-ZNF101 Antibody; Rabbit Polyclonal; ZNF101 Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-HZF12 Antibody; anti-ZNF101 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human ZNF101
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-ZNF101 antibody	IHC (Immunohistochemistry)
Immunogen	E Coli-derived Human ZNF101 fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human ZNF101 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Immunochemical staining of human ZNF101 in human testis with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunochemical staining of human ZNF101 in human liver with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.

ADDITIONAL INFORMATION

Related Product Information for anti-ZNF101 antibody	Zinc finger proteins (ZNFs), such as ZNF101, bind nucleic acids and perform many key functions, the most important of which is regulating transcription. ZNFs, which bind nucleic acids, perform many key functions, the most important of which is regulating transcription. The C2H2 ZNF family is characterized by repeated zinc finger motifs of approximately 28 amino acids that chelate a zinc ion using 2 cysteine residues and 2 histidine residues. The vast majority of these ZNFs are classified as Kruppel-like because they contain conserved 6-amino acid H/C links, the regions connecting successive finger repeats. About one-third of Kruppel-like ZNFs contain a conserved, approximately 75-amino acid motif, called the Kruppel-associated box (KRAB), in their N-terminal nonfinger region.
References	Bellefroid EJ, et al. (1993) Clustered organization of homologous krab zinc-finger genes with enhanced expression in human t lymphoid cells. Embo j 12 (4): 1363-1374.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/znf101-antibody-immunohistochemistry-elisa/256983>