

Rabbit anti-Human Bim Polyclonal Antibody | anti-BCL2L11 antibody

Anti-Bim Antibody, Rabbit Polyclonal

CATALOG # AAA257030	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

Bim; Polyclonal Antibody; Anti-Bim Antibody; Rabbit Polyclonal; Bim Antibody; Rabbit PAb; Antigen Affinity Purified; BCL2-like 11 (apoptosis facilitator); Anti-BAM Antibody; Anti-BIM Antibody; Anti-BOD Antibody; anti-BCL2L11 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human Bim
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-BCL2L11 antibody	IHC (Immunohistochemistry)
Immunogen	E Coli-derived Human Bim fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human Bim fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Immunochemical staining of human Bim in human liver with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunochemical staining of human Bim in human kidney with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.

ADDITIONAL INFORMATION

Related Product Information for anti-BCL2L11 antibody	BCL2L11, also known as Bim, belongs to the BCL-2 protein family. Members of this family form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. BCL2L11 contains a Bcl-2 homology domain 3 (BH3). It has been shown to interact with other members of the BCL-2 protein family, including BCL2, BCL2L1/BCL-X(L), and MCL1, and to act as an apoptotic activator. BCL2L11 gene functions as an essential initiator of apoptosis in thymocyte-negative selection.
Product Categories/Family for anti-BCL2L11 antibody	AKT Signaling Pathway
References	Day Catherine L. et al., 2004, Biochem J. 377 (3): 597-605.Murray S. et al., 2001, Cytogenet Cell Genet. 92 (3-4): 353.Hsu SY. et al., 1998, Mol Endocrinol. 12 (9): 1432-40.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/bim-antibody-immunohistochemistry-elisa/257030>