

Mouse anti-Human Mesothelin Monoclonal Antibody | anti-MSLN antibody

Anti-Mesothelin Antibody, Mouse Monoclonal


CATALOG # AAA257531	HOST Mouse	REACTIVITY Human	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	----------------------	----------------------------	--

PRODUCT SYNONYMS


Mesothelin; Monoclonal Antibody; Anti-Mesothelin Antibody; Mouse Monoclonal; Mesothelin/MSLN Antibody; Mouse MAb; Anti-MPF Antibody; Anti-SMRP Antibody; mesothelin; anti-MSLN antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse IgG1
Clone Number	12
Specificity	Human Mesothelin/MSLN
Purity/Purification	Protein A
Form/Format	Liquid. 0.2um filtered solution in PBS
Applicable Applications for anti-MSLN antibody	IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant Human Mesothelin/MSLN Protein
Conjugation	Unconjugated
Preparation	This antibody was produced from a hybridoma resulting from the fusion of a mouse myeloma with B cells obtained from a mouse immunized with purified, recombinant Human Mesothelin/MSLN (rh Mesothelin/MSLN; NP_005814.2; Glu296-Gly580). The IgG fraction of the cell culture supernatant was purified by Protein A affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Preservative-Free. Avoid repeated freeze-thaw cycles. This antibody is shipped as liquid solution at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.



IHC (Immunohistochemistry)
Immunochemical staining of human Mesothelin in human tonsil with mouse monoclonal antibody at 1:60 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)
Immunochemical staining of human Mesothelin in human ovarian cancer with mouse monoclonal antibody at 1:60 dilution, formalin-fixed paraffin embedded sections.

ADDITIONAL INFORMATION

Related Product Information for anti-MSLN antibody	Background: The megakaryocyte potentiating factor belongs to the mesothelin family. This family is comprised of several mammalian pre-pro-megakaryocyte potentiating factor precursor (MPF) or mesothelin proteins. Mesothelin is a glycosylphosphatidylinositol-linked glycoprotein highly expressed in mesothelial cells, mesotheliomas, and ovarian cancer, but the biological function of the protein is not known. Megakaryocyte potentiating factor is highly expressed in mesotheliomas, ovarian cancers, and some squamous cell carcinomas (at protein level). It interacts with MUC16 and potentiates megakaryocyte colony formation in vitro. Megakaryocyte potentiating factor is secreted by several mesothelioma cell lines and is frequently elevated in the blood of patients with mesothelioma. Measurement of this protein may be useful in following the response of mesothelioma to treatment.
References	Chang MC, et al. (2012) Mesothelin enhances invasion of ovarian cancer by inducing MMP-7 through MAPK/ERK and JNK pathways. <i>Biochem J.</i> 442 (2): 293-302. Nelson HH, et al. (2011) The relationship between tumor MSLN methylation and serum mesothelin (SMRP) in mesothelioma. <i>Epigenetics.</i> 6 (8): 1029-34. Bournet B, et al. (2012) Gene expression signature of advanced pancreatic ductal adenocarcinoma using low density array on endoscopic ultrasound-guided fine needle aspiration samples. <i>Pancreatology.</i> 12 (1): 27-34.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/monoclonal/mesothelin-antibody-clone-12-immunohistochemistry-elisa/257531>