

Rabbit anti-Human HDGF Polyclonal Antibody | anti-HDGF antibody

Anti-HDGF Antibody, Rabbit Polyclonal


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA257577	Rabbit	Human	Immunofluorescence, Immunocytochemistry, Immunohistochemistry

PRODUCT SYNONYMS

HDGF; Polyclonal Antibody; Anti-HDGF Antibody; Rabbit Polyclonal; HDGF Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-HMG1L2 Antibody; hepatoma-derived growth factor; anti-HDGF antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human HDGF
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid. PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-HDGF antibody	IF (Immunofluorescence), ICC (Immunocytochemistry), IHC (Immunohistochemistry)
Immunogen	E Coli-derived Human HDGF fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human HDGF fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles. This antibody is shipped as liquid solution at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.




IF (Immunofluorescence)

Immunofluorescence staining of HDGF in MCF7 cells. Cells were fixed with 4% PFA, permeabilized with 0.3% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human HDGF polyclonal antibody (dilution ratio 1:200) at 4 degree C overnight. Then cells were stained with the Alexa Fluor-488-conjugated Goat Anti-rabbit IgG secondary antibody (green). Positive staining was localized to Nucleus.



IHC (Immunohistochemistry)

Immunochemical staining of human HDGF in human liver with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunochemical staining of human HDGF in human kidney with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/hdgf-antibody-immunofluorescence-immunocytochemistry-immunohistochemistry-elisa/257577>