

Rabbit anti-Human PBDC1 Polyclonal Antibody | anti-PBDC1 antibody

Anti-PBDC1 Antibody, Rabbit Polyclonal


CATALOG # AAA257625	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunoprecipitation, Western Blot
-------------------------------	-----------------------	----------------------------	--

PRODUCT SYNONYMS


PBDC1; Polyclonal Antibody; Anti-PBDC1 Antibody; Rabbit Polyclonal; PBDC1 Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-CXorf26 Antibody; polysaccharide biosynthesis domain containing 1; anti-PBDC1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human PBDC1
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid. PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-PBDC1 antibody	IP (Immunoprecipitation), WB (Western Blot)
Immunogen	E Coli-derived Human PBDC1 fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human PBDC1 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles. This antibody is shipped as liquid solution at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.



IP (Immunoprecipitation)
CXorf26 was immunoprecipitated using: Lane A: 0.5 mg U-251 MG Whole Cell Lysate 4 uL anti-CXorf26 rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G. Primary antibody: Anti-CXorf26 rabbit polyclonal antibody, at 1:100 dilution. Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 26 kDa. Observed band size: 26 kDa.



WB (Western Blot)
Anti-CXorf26 rabbit polyclonal antibody at 1:500 dilution. Lane A: HT-1080 Whole Cell Lysate. Lane B: U-251 MG Whole Cell Lysate. Lysates/proteins at 30 ug per lane. Secondary: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 26 kDa. Observed band size: 26 kDa. (We are unsure as to the identity of these extra bands.)

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.