

Rabbit anti-Human POMGNT2 Polyclonal Antibody | anti-POMGNT2 antibody

Anti-POMGNT2 Antibody, Rabbit Polyclonal


CATALOG # AAA257632	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunoprecipitation, Western Blot
-------------------------------	-----------------------	----------------------------	--

PRODUCT SYNONYMS

POMGNT2; Polyclonal Antibody; Anti-POMGNT2 Antibody; Rabbit Polyclonal; POMGNT2/C3orf39 Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-AGO61 Antibody; Anti-GTDC2 Antibody; Anti-MDDGA8 Antibody; anti-POMGNT2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human POMGNT2/C3orf39
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid. PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-POMGNT2 antibody	IP (Immunoprecipitation), WB (Western Blot)
Immunogen	E Coli-derived Human POMGNT2/C3orf39 fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human POMGNT2/C3orf39 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles. This antibody is shipped as liquid solution at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.



IP (Immunoprecipitation)

C3orf39 was immunoprecipitated using: Lane A: 0.5 mg U-251 MG Whole Cell Lysate 4 uL anti-C3orf39 rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G. Primary antibody: Anti-C3orf39 rabbit polyclonal antibody, at 1:100 dilution Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 67 kDa Observed band size: 67 kDa



WB (Western Blot)

Anti-C3orf39 rabbit polyclonal antibody at 1:500 dilution Lane A: U-251 MG Whole Cell Lysate Lysates/proteins at 30 ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 67 kDa Observed band size: 67 kDa (We are unsure as to the identity of these extra bands).

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.