

Rabbit anti-Human OSER1 Polyclonal Antibody | anti-OSER1 antibody

Anti-OSER1 Antibody, Rabbit Polyclonal

CATALOG # AAA257633	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

OSER1; Polyclonal Antibody; Anti-OSER1 Antibody; Rabbit Polyclonal; OSER1/C20orf111 Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-C20orf111 Antibody; Anti-dJ1183I21.1 Antibody; Anti-HSPC207 Antibody; Anti-Osr1 Antibody; Anti-Perit1 Antibody; oxidative stress responsive serine-rich 1; anti-OSER1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human OSER1/C20orf111
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid. PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-OSER1 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	E Coli-derived Human OSER1/C20orf111 fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human OSER1/C20orf111 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles. This antibody is shipped as liquid solution at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

IMAGES




IHC (Immunohistochemistry)

Immunochemical staining of human C20orf111 in human liver with rabbit polyclonal antibody at 1:2000 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunochemical staining of human C20orf111 in human kidney with rabbit polyclonal antibody at 1:2000 dilution, formalin-fixed paraffin embedded sections.



WB (Western Blot)

Anti-C20orf111 rabbit polyclonal antibody at 1:500 dilution
 Lane A: Mouse liver tissue lysate
 Lysates/proteins at 30 ug per lane.
 Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.
 Developed using the ECL technique.
 Performed under reducing conditions.
 Predicted band size: 32 kDa
 Observed band size: 32 kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/oser1-antibody-immunohistochemistry-western-blot-elisa/257633>