

Rabbit anti-Human FBF1 Polyclonal Antibody | anti-FBF1 antibody

Anti-FBF1 Antibody, Rabbit Polyclonal


CATALOG # AAA257637	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunoprecipitation, Western Blot
-------------------------------	-----------------------	----------------------------	--

PRODUCT SYNONYMS

FBF1; Polyclonal Antibody; Anti-FBF1 Antibody; Rabbit Polyclonal; FBF1 Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-Alb Antibody; anti-FBF1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human FBF1
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid. PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-FBF1 antibody	IP (Immunoprecipitation), WB (Western Blot)
Immunogen	E Coli-derived Human FBF1 fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human FBF1 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles. This antibody is shipped as liquid solution at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.



IP (Immunoprecipitation)

FBF1 was immunoprecipitated using: Lane A: 0.5 mg HCT116 Whole Cell Lysate 4 uL anti-FBF1 rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G. Primary antibody: Anti-FBF1 rabbit polyclonal antibody, at 1:100 dilution. Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 125 kDa. Observed band size: 125 kDa.



WB (Western Blot)

Anti-FBF1 rabbit polyclonal antibody at 1:500 dilution. Lane A: HCT116 Whole Cell Lysate. Lane B: RT-4 Whole Cell Lysate. Lysates/proteins at 30 ug per lane. Secondary: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 125 kDa. Observed band size: 125 kDa.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.