

Rabbit anti-Human TRMT1L Polyclonal Antibody | anti-TRMT1L antibody

Anti-TRMT1L Antibody, Rabbit Polyclonal

CATALOG # AAA257641	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

TRMT1L; Polyclonal Antibody; Anti-TRMT1L Antibody; Rabbit Polyclonal; TRMT1L/C1orf25 Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-bG120K12.3 Antibody; Anti-C1orf25 Antibody; Anti-MST070 Antibody; Anti-MSTP070 Antibody; Anti-TRM1L Antibody; anti-TRMT1L antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human TRMT1L/C1orf25
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid. PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-TRMT1L antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	E Coli-derived Human TRMT1L/C1orf25 fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human TRMT1L/C1orf25 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles. This antibody is shipped as liquid solution at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

IMAGES




IHC (Immunohistochemistry)

Immunochemical staining of human C1orf25 in human adrenal gland with rabbit polyclonal antibody at 1:500 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunochemical staining of human C1orf25 in human placenta with rabbit polyclonal antibody at 1:500 dilution, formalin-fixed paraffin embedded sections.



WB (Western Blot)

Anti-C1orf25 rabbit polyclonal antibody at 1:500 dilution Lane A: Jurkat Whole Cell Lysate
Lysates/proteins at 30 ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 81 kDa Observed band size: 95 kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/trmt1l-antibody-immunohistochemistry-western-blot-elisa/257641>