

Rabbit anti-Human Bad Polyclonal Antibody | anti-Bad antibody

Anti-Bad Antibody, Rabbit Polyclonal

CATALOG # AAA257759	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

Bad; Polyclonal Antibody; Anti-Bad Antibody; Rabbit Polyclonal; Bad Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-BBC2 Antibody; Anti-BCL2L8 Antibody; BCL2-associated agonist of cell death; anti-Bad antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human Bad
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid. PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-Bad antibody	IHC (Immunohistochemistry)
Immunogen	E Coli-derived Human Bad fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human Bad fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles. This antibody is shipped as liquid solution at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

IMAGES



IHC (Immunohistochemistry)

Immunochemical staining of human Bad in human kidney with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunochemical staining of human Bad in human kidney with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.

ADDITIONAL INFORMATION

Related Product Information for anti-Bad antibody	Background: Bcl-2 agonist of cell death (BAD) is a BH3-only member of the Bcl-2 family which possesses important regulatory function in apoptosis. BAD has also been shown to possess many non-apoptotic functions closely linked to cancer including regulation of glycolysis, autophagy, cell cycle progression and immune system development. Oral squamous cell carcinoma (OSCC) is a type of cancer with high morbidity and mortality rates worldwide; it also demonstrates chemotherapeutic resistance. Triterpenoid ursolic acid has been shown to exhibit various biological activities and anticancer effects in several preclinical studies. The apoptosis-associated signaling showed that ursolic acid decreased the phosphorylation of AKT (Ser473) and B-cell lymphoma 2 (Bcl-2)-associated agonist of cell death (BAD; Ser136), and the protein levels of Bcl-2 and Bcl-extra large (Bcl-xL), and increased the expression of BAD and Bcl-2-associated X (Bax) protein in CAR cells.
Product Categories/Family for anti-Bad antibody	TNF Signaling

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/bad-antibody-immunohistochemistry-elisa/257759>