

EPOR active protein

EPOR Protein, Human, Recombinant (ECD, His Tag)

CATALOG #
AAA257919

HOST
HEK293 Cells



PRODUCT SYNONYMS

EPOR; EPOR Protein; Human; Recombinant (ECD; His Tag); Human EPO Receptor/EPOR Protein (ECD; EPO-R Protein; erythropoietin receptor; EPOR active protein

PRODUCT SPECIFICATIONS

Host	HEK293 Cells
Purity/Purification	>98% as determined by SDS-PAGE
Form/Format	Lyophilized from sterile PBS, 0.05% CHAPS, pH7.4. Normally 5%-8% trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the hard copy of CoA.
Sequence	Met1-Pro250
Species	Human
Activity	Measured by its ability to inhibit EPO-dependent proliferation of TF-1 human erythroleukemic cells. The ED50 for this effect is typically 15-60ng/mL in the presence of 0.1 U/mL Recombinant Human EPO.
Endotoxin	<1.0EU per ug of the protein as determined by the LAL method
Predicted N Terminal	Ala 25
Tag	C-His
Protein Construction	A DNA sequence encoding the extracellular domain (Met 1-Pro 250) of human erythropoietin receptor (NP_000112.1) precursor was fused with a polyhistidine tag at the C-terminus.
Reconstitution	A hardcopy of COA with reconstitution instruction is sent along with the products. Please refer to it for detailed information.
Preparation and Storage	Samples are stable for up to twelve months from date of receipt at -20 degree C to -80 degree C. Store it under sterile conditions at -20 degree C to -80 degree C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles. In general, recombinant proteins are provided as lyophilized powder which are shipped at ambient temperature. Bulk packages of recombinant proteins are provided as frozen liquid. They are shipped out with blue ice.

IMAGES



SDS-PAGE

Application Data
Measured by its ability to inhibit EPO-dependent proliferation of TF-1 human erythroleukemic cells. The ED50 for this effect is typically 15-60 ng/mL in the presence of 0.1 U/mL Recombinant Human EPO.

ADDITIONAL INFORMATION

Related Product Information for EPOR active protein	Background: Erythropoietin (EPO) is the major glycoprotein hormone regulator of mammalian erythropoiesis, and is produced by kidney and liver in an oxygen-dependent manner. The biological effects of EPO are mediated by the specific erythropoietin receptor (EPOR/EPO Receptor) on bone marrow erythroblasts, which transmits signals important for both proliferation and differentiation along the erythroid lineage. EPOR protein is a type α- single-transmembrane cytokine receptor, and belongs to the homodimerizing subclass which functions as ligand-induced or ligand-stabilized homodimers. EPOR signaling prevents neuronal death and ischemic injury. Recent studies have shown that EPO and EPOR protein may be involved in carcinogenesis, angiogenesis, and invasion.
References	Divoky V, et al. (2002) Mouse surviving solely on human erythropoietin receptor (EpoR): model of human EpoR-linked disease. Blood 99(10): 3873-4.Carruthers SG. (2009) A truncated erythropoietin receptor EPOR-T is associated with hypertension susceptibility. Clin Pharmacol Ther. 86(2): 134-6.Baltaziak M, et al. (2009) Relationships of P53 and Bak with EPO and EPOR in human colorectal cancer. Anticancer Res. 29(10):4151-6.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4503591
NCBI GeneID	2057
NCBI Accession #	NP_000112.1
NCBI GenBank Nucleotide #	NM_000121.3
UniProt Accession #	P19235; Q15443; Q2M205; B2RCG4
Molecular Weight	55,065 Da
NCBI Official Full Name	erythropoietin receptor
NCBI Official Synonym Full Names	erythropoietin receptor
NCBI Official Symbol	EPOR
NCBI Official Synonym Symbols	EPO-R
NCBI Protein Information	erythropoietin receptor
UniProt Protein Name	Erythropoietin receptor
UniProt Gene Name	EPOR
UniProt Synonym Gene Names	EPO-R

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/hek293-active-protein/epor/257919>