

Follistatin Active Protein | FST active protein

Follistatin Protein, Human, Recombinant (His Tag)

CATALOG #
AAA258012



HOST
HEK293 Cells

PRODUCT SYNONYMS

Follistatin; Follistatin Protein; Human; Recombinant (His Tag); Human Follistatin/FST (FS288) Protein (His Tag); FS Protein; follistatin; FST active protein

PRODUCT SPECIFICATIONS

Host	HEK293 Cells
Purity/Purification	>90% as determined by SDS-PAGE
Form/Format	Lyophilized from sterile PBS, pH7.4. Normally 5%-8% trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the hard copy of CoA.
Sequence	Met1-Asn31 7
Species	Human
Activity	Measured by its ability to neutralize Activin-mediated inhibition on MPC11 cell proliferation. The ED50 for this effect is typically 20-80ng/mL in the presence of 10ng/mL Recombinant Human Activin A.
Endotoxin	<1.0EU per ug of the protein as determined by the LAL method
Predicted N Terminal	Gly 30
Tag	C-His
Protein Construction	A DNA sequence encoding the human FST isoform FST288 (P19883-2)(Met 1-Asn 317) was fused with a polyhistidine tag at the C-terminus.
Reconstitution	A hardcopy of COA with reconstitution instruction is sent along with the products. Please refer to it for detailed information.
Preparation and Storage	Samples are stable for up to twelve months from date of receipt at -20 degree C to -80 degree C. Store it under sterile conditions at -20 degree C to -80 degree C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles. In general, recombinant proteins are provided as lyophilized powder which are shipped at ambient temperature. Bulk packages of recombinant proteins are provided as frozen liquid. They are shipped out with blue ice.



SDS-PAGE

Application Data
Measured by its ability to neutralize Activin-mediated inhibition on MPC11 cell proliferation.
The ED50 for this effect is typically 20-80 ng/mL in the presence of 10 ng/mL Recombinant Human Activin A.

ADDITIONAL INFORMATION

Related Product Information for FST active protein	Background: Follistatin is a single-chain gonadal protein that specifically inhibits follicle-stimulating hormone release. The single FST gene encodes two isoforms, FST317 and FST344 containing 317 and 344 amino acids respectively, resulting from alternative splicing of the precursor mRNA. In a study in which 37 candidate genes were tested for linkage and association with polycystic ovary syndrome (PCOS) or hyperandrogenemia in 150 families, evidence was found for linkage between PCOS and follistatin. follistatin are expressed and subserve local regulatory roles in numerous extragonadal tissues, including brain, adrenal, bone marrow, and placenta but perhaps most notably in anterior pituitary-the classical target tissue for inhibin, the activin-follistatin system may play a key role in early embryogenesis. Follistatin binds directly to activin and functions as an activin antagonist. Specific inhibitor of the biosynthesis and secretion of pituitary follicle stimulating hormone follistatin is a binding protein to activin. Since activin binds to follistatin, it is imperative to determine the nature of the activin/follistatin binding complex.
References	Carroll RS, et al. (1989) Inhibin, activin, and follistatin: regulation of follicle-stimulating hormone messenger ribonucleic acid levels. Mol Endocrinol. 3(12): 1969-76.Fainsod A, et al. (1997) The dorsalizing and neural inducing gene follistatin is an antagonist of BMP-4. Mech Dev. 63(1): 39-50.Kaiser UB, et al. (1992) Follistatin gene expression in the pituitary: localization in gonadotropes and folliculostellate cells in diestrous rats. Endocrinology. 130(5): 3048-56.

NCBI AND UNIPROT INFORMATION

NCBI GI #	5453652
NCBI GeneID	10468
NCBI Accession #	NP_006341.1
NCBI GenBank Nucleotide #	NM_006350.3
UniProt Accession #	P19883; Q9BTH0; B5BU94
Molecular Weight	38,007 Da
NCBI Official Full Name	follistatin isoform FST317
NCBI Official Synonym Full Names	follistatin
NCBI Official Symbol	FST
NCBI Official Synonym Symbols	FS
NCBI Protein Information	follistatin; activin-binding protein; follistatin isoform FST317
UniProt Protein Name	Follistatin

UniProt Gene Name	FST
UniProt Synonym Gene Names	FS
UniProt Entry Name	FST_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/hek293-fst-active-protein/follistatin/258012>