

Rabbit anti-Human HNRNPK Polyclonal Antibody | anti-HNRNPK antibody

HNRNPK Antibody, Rabbit PAb, Antigen Affinity Purified

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA258658	Rabbit	Human	Immunoprecipitation, Immunofluorescence, Immunocytochemistry, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

HNRNPK; Polyclonal Antibody; HNRNPK Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-AUKS Antibody; Anti-CSBP Antibody; Anti-HNRPK Antibody; Anti-TUNP Antibody; anti-HNRNPK antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	Human HNRNPK
Applicable Applications for anti-HNRNPK antibody	IP (Immunoprecipitation), IF (Immunofluorescence), ICC (Immunocytochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Preparation	Produced in rabbits immunized with E Coli-derived Human HNRNPK fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 to -80 degree C. Preservative-Free. Avoid repeated freeze-thaw cycles.

IMAGES





IP (Immunoprecipitation)

HNRNPK was immunoprecipitated using:; Lane A:0.5 mg HeLa Whole Cell Lysate; Lane B:0.5 mg Jurkat Whole Cell Lysate; Lane C:0.5 mg A431 Whole Cell Lysate; Lane D:0.5 mg HepG2 Whole Cell Lysate; 4 uL anti-HNRNPK rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G.; Primary antibody:; Anti-HNRNPK rabbit polyclonal antibody, at 1:100 dilution; Secondary antibody:; Clean-Blot IP Detection Reagent (HRP) at 1:1000 dilution; Developed using the ECL technique.; Performed under reducing conditions.; Predicted band size: 60 kDa; Observed band size :60 kDa




IF (Immunofluorescence)

Immunofluorescence staining of HNRNPK in HeLa cells. Cells were fixed with 4% PFA, permeabilized with 0.3% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human HNRNPK polyclonal antibody (dilution ratio 1:100) at 4 degree C overnight. Then cells were stained with the Alexa Fluor 488-conjugated Goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI (blue). Positive staining was localized to Nucleus.



IHC (Immunohistochemistry)
 Immunochemical staining of human HNRNPK in human skin with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.

IHC (Immunohistochemistry)
 Immunochemical staining of human HNRNPK in human lymph node with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



WB (Western Blot)
 Anti-HNRNPK rabbit polyclonal antibody at 1:500 dilution; Lane A: HeLa Whole Cell Lysate; Lane B: Jurkat Whole Cell Lysate; Lane C: A431 Whole Cell Lysate; Lane D: HepG2 Whole Cell Lysate; Lysates/proteins at 30 ug per lane.; Secondary: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.; Developed using the ECL technique.; Performed under reducing conditions.; Predicted band size: 51 kDa; Observed band size: 60 kDa

ADDITIONAL INFORMATION

Product Categories/Family for anti-HNRNPK antibody	Epigenetics
--	-------------

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://www.aaabiotech.com/polyclonal/hnrnpk-antibody-immunoprecipitation-immunofluorescence-immunocytochemistry-immunohistochemistry-western-blot-elisa/258658>