

Rabbit anti-Human CD79A Polyclonal Antibody | anti-CD79A antibody

Anti-CD79A Antibody, Rabbit Polyclonal


CATALOG # AAA258711	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunoprecipitation, Western Blot
-------------------------------	-----------------------	----------------------------	--

PRODUCT SYNONYMS

CD79A; Polyclonal Antibody; Anti-CD79A Antibody; Rabbit Polyclonal; CD79a molecule; immunoglobulin-associated alpha; Anti-IGA Antibody; Anti-MB-1 Antibody; anti-CD79A antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human CD79A
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; 0.2um filtered solution in PBS
Applicable Applications for anti-CD79A antibody	IP (Immunoprecipitation), WB (Western Blot)
Immunogen	A synthetic peptide corresponding to the C-terminus of the Human CD79A
Preparation	Produced in rabbits immunized with a synthetic peptide corresponding to the C-terminus of the Human CD79A, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Preservative-Free. Avoid repeated freeze-thaw cycles.



IP (Immunoprecipitation)

Lane A: 0.5 mg Daudi Whole Cell Lysate 4 uL anti-CD79a rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G. Primary antibody: Anti-CD79a rabbit polyclonal antibody, at 1:100 dilution Secondary antibody: Clean-Blot IP Detection Reagent (HRP) at 1:1000 dilution Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 25 kDa Observed band size : 40-55 kDa



WB (Western Blot)

Lane A: Daudi whole cell lysate Lysates/proteins at 30 ug per lane.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.